arXiv:1210.8316v2 [math.AG] 3 Nov 2012

The number of singular vector tuples and uniqueness of best rank one approximation of tensors

Shmuel Friedland* Giorgio Ottaviani†

Abstract

In this paper we discuss the notion of singular value tuples of a complex valued d-mode tensor of dimension $m_1 \times \ldots \times m_d$. We show that a general tensor has a finite number of singular value tuples, viewed as points in a corresponding Segre variety. We give the formula for the number of singular value tuples. We show similar results for tensors with partial symmetry. We give analogous results for the homogeneous pencil eigenvalue problem for cubic tensors, i.e. $m_1 = \ldots = m_d$. We show that best rank one approximations for general tensors are unique. Similarly, for general partially symmetric tensors the best rank one approximation is unique and partially symmetric. We show that a best rank-$(r_1, \ldots, r_d)$ approximation for a general d-mode tensor is unique.

2010 Mathematics Subject Classification. 14D21, 15A18, 15A69, 65D15, 65H10, 65K05.

Key words. Singular value tuples, vector bundles, Chern classes, partially symmetric tensors, homogeneous pencil eigenvalue problem for cubic tensors, Singular value decomposition, rank one approximation, a best rank-$(r_1, \ldots, r_d)$ approximation.

1 Introduction

The object of this paper is to study two closely related topics: counting the number of singular vector tuples of a d-mode complex generic tensor of dimension $m_1 \times \ldots \times m_d$ and the uniqueness of its best rank one approximation. To motivate our results let us consider the classical case of matrices, i.e. $d = 2$ and $A \in \mathbb{R}^{m_1 \times m_2} \equiv \mathbb{R}^{m_1} \otimes \mathbb{R}^{m_2}$. We call a pair $(\mathbf{x}_1, \mathbf{x}_2) \in (\mathbb{R}^{m_1} \setminus \{\mathbf{0}\}) \times (\mathbb{R}^{m_2} \setminus \{\mathbf{0}\})$ a *singular vector pair* if

$$A\mathbf{x}_2 = \lambda_1 \mathbf{x}_1, \quad A^\top \mathbf{x}_1 = \lambda_2 \mathbf{x}_2, \tag{1.1}$$

for some $\lambda_1, \lambda_2 \in \mathbb{R}$. For $\mathbf{x} \in \mathbb{R}^m$ let $\|\mathbf{x}\| := \sqrt{\mathbf{x}^\top \mathbf{x}}$ be the Euclidean norm on $\mathbb{R}^m$. Choosing $\mathbf{x}_1, \mathbf{x}_2$ to be of Euclidean length one we deduce that $\lambda_1 = \lambda_2$, where $|\lambda_1|$ is equal to some singular value of A. It is natural to identify all singular vector pairs of the form $(a_1\mathbf{x}_1, a_2\mathbf{x}_2)$, where $a_1 a_2 \neq 0$ as the class of singular vector pair. This is equivalent to view a singular vector pair as an element of $\mathbb{P}(\mathbb{R}^{m_1}) \times \mathbb{P}(\mathbb{R}^{m_2})$, where $\mathbb{P}(\mathbb{F}^m)$ is the space of all lines through the origin in $\mathbb{F}^m$ for any field $\mathbb{F}$. For a generic A, i.e. A of the maximal rank $r = \min(m_1, m_2)$ and r distinct positive

---

[*] Dept. of Mathematics, Statistics and Computer Science, Univ. of Illinois at Chicago, Chicago, Illinois 60607-7045, USA, `friedlan@uic.edu`. Supported by NSF grant DMS–1216393.

[†] Dipartimento di Matematica e Informatica "Ulisse Dini", Università di Firenze, viale Morgagni 67/A, 50134 Firenze, Italy, `ottavian@math.unifi.it`. Member of INDAM.

singular values, A has exactly r distinct singular vector pairs. Furthermore, under these conditions A has a unique best rank one approximation in the Frobenius norm given by the singular pair corresponding to the maximal singular value [8].

Assume now that $m = m_1 = m_2$ and A is a real symmetric matrix. Then the singular values of A are the absolute values of the eigenvalues of A. Furthermore, if all the absolute values of the eigenvalues of A are pairwise distinct then A has a unique best rank one approximation which is symmetric. Hence for any real symmetric matrix A there exists a best rank one approximation which is symmetric.

In this paper we derive similar results for tensors. To state more precisely our results we introduce some notation. Let $\mathbb{F}$ be either the real field $\mathbb{R}$ or the complex field $\mathbb{C}$. Let $m_i \geq 2$ be an integer for $i \in [d] := \{1, \ldots, d\}$. Denote $\mathbf{m} := (m_1, \ldots, m_d)$. Let $\Sigma_{\mathbb{F}}(\mathbf{m}) := \mathbb{P}(\mathbb{F}^{m_1}) \times \ldots \times \mathbb{P}(\mathbb{F}^{m_d})$ be the Segre variety over $\mathbb{F}$. Set $\Sigma(\mathbf{m}) := \Sigma_{\mathbb{C}}(\mathbf{m})$.

Denote by $\mathbb{F}^{\mathbf{m}} = \mathbb{F}^{m_1 \times \ldots \times m_d} := \otimes_{i=1}^d \mathbb{F}^{m_i}$ the vector space of d-mode tensors $\mathcal{T} = [t_{i_1,\ldots,i_d}], i_j = 1, \ldots, m_j, j = 1, \ldots, d$ over $\mathbb{F}$. (We assume that $d \geq 3$ unless stated otherwise.) For an integer $p \in [d]$ and for $\mathbf{x}_{j_r} \in \mathbb{F}^{m_{j_r}}, r \in [p]$ we use the notation $\otimes_{j_r, r\in[p]} \mathbf{x}_{j_r} := \mathbf{x}_{j_1} \otimes \ldots \otimes \mathbf{x}_{j_p}$. For a subset $P = \{j_1, \ldots, j_p\} \subseteq [d]$ of cardinality $p = |P|$, consider a p-mode tensor $\mathcal{X} = [x_{i_{j_1},\ldots,i_{j_p}}] \in \otimes_{j_r, r\in[p]} \mathbb{F}^{m_{j_r}}$, where $j_1 < \ldots < j_p$. Define $\mathcal{T} \times \mathcal{X} := \sum_{i_{j_r} \in [m_{j_r}], r\in[p]} t_{i_1,\ldots,i_d} x_{i_{j_1},\ldots,i_{j_p}}$ to be a $(d-p)$-mode tensor obtained by contraction on the indices $i_{j_1}, \ldots, i_{j_p}$.

Let $\mathcal{T} \in \mathbb{F}^{\mathbf{m}}$. We first define the notion of a singular vector tuple $(\mathbf{x}_1, \ldots, \mathbf{x}_d) \in (\mathbb{F}^{m_1} \setminus \{\mathbf{0}\}) \times \ldots \times (\mathbb{F}^{m_d} \setminus \{\mathbf{0}\})$ [14]:

$$\mathcal{T} \times \otimes_{j\in[d]\setminus\{i\}} \mathbf{x}_j = \lambda_i \mathbf{x}_i, \quad i = 1, \ldots, d. \tag{1.2}$$

As for matrices we identify all singular vector tuples of the form $(a_1\mathbf{x}_1, \ldots, a_d\mathbf{x}_d)$, $a_1 \ldots a_d \neq 0$ as one class of singular vector tuple in $\Sigma_{\mathbb{F}}(\mathbf{m})$. (Note that for $d = 2$ and $\mathbb{F} = \mathbb{C}$ our notion of singular pair differs from the classical notion of singular vectors for complex valued matrices.) We remark that, as in the classical SVD, $\mathbb{F}^{m_i}$ cannot be replaced by a vector space V_i of dimension m_i, unless we choose a basis which allows to identify V_i with its dual $V_i^\vee$. Our first major result is that for a generic tensor $\mathcal{T} \in \mathbb{C}^{\mathbf{m}}$ the number of singular vectors tuples in $(\mathbf{x}_1, \ldots, \mathbf{x}_d) \in \Sigma(\mathbf{m})$ is finite and it is equal to $c(\mathbf{m})$. $c(\mathbf{m})$ is the coefficient of the monomial $\prod_{i\in[d]} t_i^{m_i-1}$ in the homogenous polynomial in variables $t_1, \ldots, t_d$

$$\prod_{i\in[d]} \frac{\hat{t}_i^{m_i} - t_i^{m_i}}{\hat{t}_i - t_i}, \quad \hat{t}_i = \sum_{j\in[d]\setminus\{i\}} t_j, \ i \in [d]. \tag{1.3}$$

In particular, for a generic $\mathcal{T} \in \mathbb{R}^{m_1\times\ldots\times m_d}$ the number of singular vectors tuples in $(\mathbf{x}_1, \ldots, \mathbf{x}_d) \in \mathbb{P}(\mathbb{R}^{m_1}) \times \ldots \times \mathbb{P}(\mathbb{R}^{m_d})$ is finite and it is at most $c(\mathbf{m})$. At the end of §3 we list the first values of $c(\mathbf{m})$ for $d = 3$. We generalize the above results for the class of tensors with given partial symmetry.

We now consider the cubic case where $m_1 = \ldots = m_d = m$. For an integer $m \geq 2$ let $m^{\times d} := (\underbrace{m, \ldots, m}_{d})$. Then $\mathcal{T} \in \mathbb{F}^{m^{\times d}}$ is called d-cube, or simply a cube tensor. For a vector $\mathbf{x} \in \mathbb{C}^m$ let $\otimes^k \mathbf{x} := \underbrace{\mathbf{x} \otimes \ldots \otimes \mathbf{x}}_{k}$. Assume that $\mathcal{T}, \mathcal{S} \in \mathbb{C}^{m^{\times d}}$. Then the homogeneous pencil eigenvalue problem is $\mathcal{T} \times \otimes^{d-1}\mathbf{x} = \lambda \mathcal{S} \times \otimes^{d-1}\mathbf{x}$. The contraction here is with respect to the last $d-1$ indices of $\mathcal{T},\mathcal{S}$ respectively. We assume without the loss of generality that $\mathcal{T} = [t_{i_1,\ldots,i_d}], \mathcal{S} = [s_{i_1,\ldots,i_d}]$ are symmetric with respect to the indices $i_2, \ldots, i_d$. $\mathcal{S}$ is called nonsingular if the system $\mathcal{S} \times \otimes^{d-1}\mathbf{x} = \mathbf{0}$ implies that $\mathbf{x} = 0$. Assume that $\mathcal{S}$ is nonsingular and fixed. Then $\mathcal{T}$ has exactly $m(d-1)^{m-1}$ eigenvalues counted with their multiplicities. $\mathcal{T}$ has $m(d-1)^{m-1}$ distinct eigenvectors eigenvectors in $\mathbb{P}(\mathbb{C}^m)$ for a general $\mathcal{T}$. See [19] for the case $\mathcal{S}$ is the identity tensor.

View $\mathbb{R}^{m_1\times\ldots m_d}$ as an inner product space, where for two d-mode tensors $\mathcal{T},\mathcal{S} \in \mathbb{R}^{m_1\times\ldots\times m_d}$ we let $\langle \mathcal{T},\mathcal{S}\rangle := \mathcal{T} \times \mathcal{S}$. Then the Hilbert-Schmidt norm is defined $\|\mathcal{T}\| := \sqrt{\langle \mathcal{T},\mathcal{T}\rangle}$. (For simplicity of notation, we call the Frobenius and the Hilbert-Schmidt norm L_2 norms.) The best rank one approximation is a solution to the minimal problem

$$\min_{\mathbf{x}_i\in\mathbb{R}^{m_i}, i\in[d]} \|\mathcal{T} - \otimes_{i\in[d]}\mathbf{x}_i\| = \|\mathcal{T} - \otimes_{i\in[d]}\mathbf{u}_i\|. \tag{1.4}$$

$\otimes_{i\in[d]}\mathbf{u}_i$ is called best rank one approximation of $\mathcal{T}$. In this paper we show that for a generic $\mathcal{T} \in \mathbb{R}^{m_1\times\ldots\times m_d}$ the best rank one approximation is unique.

We generalize the above results for the class of tensors with given partial symmetry. In particular we show that a generic real symmetric tensor $\mathcal{T}$ has a unique rank one approximation which satisfy (1.4). Hence this unique rank one tensor is symmetric. This gives another proof that the best rank one approximation to a symmetric tensor can be chosen symmetric [5]. (See a related result [21].)

In the last section of this paper we study a best rank-$(r_1,\ldots,r_d)$ approximation for a d-mode tensor [4]. We show that for a general tensor a best rank-$(r_1,\ldots,r_d)$ approximation is unique.

We now describe briefly the contents of our paper. In §2 we give layman's introduction to some basic notions of vector bundles over compact complex manifolds and Chern classes of certain bundles over the Segre varieties needed for this paper. We hope that this introduction will make our paper accessible to a wider audience. §3 discusses the first main contribution of this paper. Namely, the number of singular value tuples of a general complex tensor is finite and is equal to $c(m_1,\ldots,m_d)$. We give a closed formula for $c(m_1,\ldots,m_d)$. §4 generalizes these results to partially symmetric tensors. In particular we reproduce the result of Cartwright and Sturmfels for symmetric tensors [2]. In §5 we discuss a homogeneous pencil eigenvalue problem. In §6 we give certain conditions on a very general best approximation problem in $\mathbb{R}^n$, which are probably well known to the experts. §7 give uniqueness results on best rank one approximation of general partially symmetric tensors. In §8 we discuss a best rank-$(r_1,\ldots,r_d)$ approximation.

2 Vector bundles over compact complex manifolds

In this section we recall some basic results on complex manifolds and holomorphic tangents bundles that we use in this paper. Our object was to give the simplest possible intuitive description of basic results in algebraic geometry needed in this paper, sometimes compromising the rigor. An interested reader can consult for more details with [9] for general facts about complex manifolds and complex vector bundles, and for a simple axiomatic exposition on complex vector bundles with [13]. For Bertini's type theorem we refer to Fulton [6] and Hartshorne [10].

2.1 Complex compact manifolds

Let M be a compact complex manifold of dimension n. Thus there exists a finite open cover $\{U_i\}, i \in [N]$ with coordinate homeomorphism $\phi_i : U_i \to \mathbb{C}^n$ such that $\phi_i \circ \phi_j^{-1}$ is holomorphic $\phi_j(U_i \cap U_j)$ for all i, j.

As an example consider the $m-1$ dimensional complex projective space $\mathbb{P}(\mathbb{C}^m)$, which is the set of all complex lines in $\mathbb{C}^m$ through the origin. Any point in $\mathbb{P}(\mathbb{C}^m)$ is represented by a one dimensional subspace spanned by the vector $\mathbf{x} = (x_1, \ldots, x_m)^\top \in \mathbb{C}^m \setminus \{\mathbf{0}\}$. The standard open cover of $\mathbb{P}(\mathbb{C}^m)$ consists of m open covers $U_1, \ldots, U_m$, where U_i corresponds to the lines spanned by $\mathbf{x}$ with $x_i \neq 0$. The homeomorphism ϕ_i is given by $\phi_i(\mathbf{x}) = (\frac{x_1}{x_i}, \ldots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \ldots, \frac{x_m}{x_i})^\top$. So each U_i is homeomorphic to $\mathbb{C}^{m-1}$.

Let M be an n-dimensional compact complex manifold as above. For $\zeta \in U_i$, the coordinates of the vector $\phi_i(\zeta) = \mathbf{z} = (z_1, \ldots, z_n)^\top$ are called the local coordinates of ζ. Since $\mathbb{C}^n \equiv \mathbb{R}^{2n}$, M is a real manifold of real dimension $2n$. Let $z_j = x_j + \mathbf{i}y_j, \bar{z}_j = x_j - \mathbf{i}y_j, j \in [n]$, where $\mathbf{i} = \sqrt{-1}$. For simplicity of notation we let $\mathbf{u} = (u_1, \ldots, u_{2n}) = (x_1, y_1, \ldots, x_n, y_n)$ be the real local coordinates on U_i. Any function $f : U_i \to \mathbb{C}$ in the local coordinates is viewed as $f(\mathbf{u}) = g(\mathbf{u}) + \mathbf{i}h(\mathbf{u})$, where $h, g : U_i \to \mathbb{R}$. Thus $df = \sum_{j \in [2n]} \frac{\partial f}{\partial u_j} du_j$. For a positive integer p, a (differential) p-form ω on U_i is given in the local coordinates as follows

$$\omega = \sum_{1 \le i_1 < \ldots < i_p \le 2n} f_{i_1, \ldots, i_p}(\mathbf{u}) du_{i_1} \wedge \ldots \wedge du_{i_p}.$$

Recall that the wedge product of two differential is anti commutative. i.e. $du_k \wedge du_l = -du_l \wedge du_k$. Then

$$d\omega = \sum_{1 \le i_1 < \ldots < i_p \le 2n} (df_{i_1, \ldots, i_p}) \wedge du_{i_1} \wedge \ldots \wedge du_{i_p}.$$

(0-differential form is a function.) Note that for $p > 2n$ any p differential form is a zero form. A straightforward calculation shows that $d(d\omega) = 0$. ω is a p form on M if its restriction to each U_i is an p form, and the restrictions of these two forms on $U_i \cap U_j$ are obtained one from the other one by the change of coordinates $\phi_i \circ \phi_j^{-1}$. ω is called closed if $d\omega = 0$, and $d\omega$ is called an exact form. The space of closed p forms modulo exact p forms is a finite dimensional vector space over $\mathbb{C}$, which is denoted by $\mathrm{H}^p(M)$. Each element of $\mathrm{H}^p(M)$ is represented by a p closed form, and the difference between two representatives is an exact form. Since the product of two forms is also a form, it follows that all closed forms modulo exact forms is a finite dimensional algebra, where the identity 1 corresponds to constant function with value 1 on M.

2.2 Holomorphic vector bundles

A holomorphic vector bundle E on M of rank k, where k is a nonnegative integer, is a complex manifold of dimension $n + k$ which can be simply described as follows. There exists a finite open cover $\{U_i\}, i \in [N]$ of M with the properties as above satisfying the following additional conditions. At each $\zeta \in U_i$ we are given k-dimensional vector space E_ζ, called a fiber of E over ζ, which all can be identified with a fixed vector space $\mathbf{V}_i$, having a basis $[\mathbf{e}_{1,i}, \ldots, \mathbf{e}_{k,i}]$. For $\zeta \in U_i \cap U_j, i \neq j$ the transition matrix from $[\mathbf{e}_{1,i},\ldots,\mathbf{e}_{k,i}]$ to $[\mathbf{e}_{1,j},\ldots,\mathbf{e}_{k,j}]$ is given by an $k\times k$ invertible matrix $g_{U_jU_i}(\zeta)$. So $[\mathbf{e}_{1,i},\ldots,\mathbf{e}_{k,i}]=[\mathbf{e}_{1,j},\ldots,\mathbf{e}_{k,j}]g_{U_jU_i}(\zeta)$. Each entry of $g_{U_jU_i}(\zeta)$ is a holomorphic function in the local coordinates of U_j. We have the following relations

$$g_{U_iU_j}(\zeta)g_{U_jU_i}(\zeta)=g_{U_iU_j}(\eta)g_{U_jU_p}(\eta)g_{U_pU_i}(\eta)=I_k \text{ for } \zeta\in U_i\cap U_j, \eta\in U_i\cap U_j\cap U_p.$$

(I_k is an identity matrix of order k.)

For $k=0$ E is called a zero bundle. E is called a line bundle if $k=1$. E is called a trivial bundle if there exists a finite open cover such that each $g_{U_iU_j}(\zeta)$ is an identity matrix. A vector bundle F on M is called a subbundle of E if F is a submanifold of E such that F_ζ is a subspace of E_ζ for each $\zeta\in M$. Assume that F is a subbundle of E. Then $G:=E/F$ is the quotient bundle of E and F, where G_ζ is the quotient vector space E_ζ/F_ζ. Let E_1,E_2 be two vectors bundles on M. We can create the following new bundles on M. $E:=E_1\oplus E_2, F:=E_1\otimes E_2$ where $E_\zeta=E_{1,\zeta}\oplus E_{2,\zeta}, F_\zeta=E_{1,\zeta}\otimes E_{2,\zeta}$. In particular, for a given vector bundle E on M we can define the bundle $F:=\otimes^d E$. Here $F_\zeta=\otimes^d E_\zeta$ is a fiber of d-mode tensors.

We now discuss a basic example used in this paper. Consider the trivial bundle E on $\mathbb{P}(\mathbb{C}^m)$ of rank m. So $E_\zeta=\mathbb{C}^m$, and by abuse of notation we denote E as $\mathbb{C}^m$. The tautological line bundle $T(m)$ on $\mathbb{P}(\mathbb{C}^m)$, customary denoted by $\mathcal{O}(-1)$, is given by $T(m)_{[\mathbf{x}]}=\mathrm{span}(\mathbf{x})\subset\mathbb{C}^m$, where $\mathbf{x}$ is a nonzero vector spanning the line through zero that represents a point $[\mathbf{x}]$ in $\mathbb{P}(\mathbb{C}^m)$. So $T(m)$ is a subbundle of $\mathbb{C}^m$. Denote by $Q(m)$ the quotient bundle $\mathbb{C}^m/T(m)$. So rank $Q(m)=m-1$. We have an exact sequence of the following bundles on $\mathbb{P}(\mathbb{C}^m)$

$$0\to T(m)\to\mathbb{C}^m\to Q(m)\to 0. \tag{2.1}$$

The dual of the subbundle $T(m)$ of $\mathbb{C}^m$, also called the hyperplane line bundle, denoted here by $H(m)$, associates with a point $[\mathbf{x}]\in\mathbb{P}(\mathbb{C}^m)$ the hyperplane $\mathbf{y}\in\mathbb{C}^{m\vee},\mathbf{x}^\top\mathbf{y}=0$ (let's remind that the initial choice of the standard basis identifies $\mathbb{C}^m$ with its dual). $H(m)$ is customary denoted by $\mathcal{O}(1)$.

2.3 Chern polynomials

We now return to a holomorphic vector bundle E on a compact complex manifold M. One associate with E the Chern polynomial $C(t,E)=1+\sum_{j=1}^{\mathrm{rank}\,E}c_j(E)t^j$ (see [9, 12]). Here $c_j(E)\in\mathrm{H}^{2j}(M)$. Note that $c_j(E)=0$ for $j>2n$. $C(t,E)=1$ if E is a trivial line bundle. $C(t,E_1\oplus E_2)=C(t,E_1)C(t,E_2)$, (Whitney sum formula). Furthermore, for a pull back bundle $\pi_1^*E_1$ described above we have the equality $c_j(\pi_1^*E_1)=c_j(E_1)$, when we use the local coordinates $\zeta=(\zeta_1,\zeta_2)$ on $M_1\times M_2$.

We now summarize the information on Chern classes needed in this paper. First, on $\mathbb{P}(\mathbb{C}^m)$ Chern classes may be identified with integers, and we can write $C(t,T(m))=1-t, C(t,H(m))=1+t$. Moreover the algebra of all closed forms modulo the exact forms on $\mathbb{P}(\mathbb{C}^m)$ is $\mathbb{C}[t]/(t^m)$, i.e. all polynomials in the variable t modulo the relation $t^m=0$. The exact sequence (2.1) yields that $C(t,Q(m))=\frac{1}{1-t}=1+\sum_{j=1}^{m-1}t^j$, that is $c_i(Q(m))=1$ for $i=1,\ldots,m-1$.

2.4 Certain bundles on Segre varieties

Let $m_1,\ldots,m_d\geq 2$ be given integers with $d>1$ and denote $\mathbf{m}=(m_1,\ldots,m_d)$. Consider the Segre variety $\Sigma(\mathbf{m}):=\mathbb{P}(\mathbb{C}^{m_1})\times\ldots\times\mathbb{P}(\mathbb{C}^{m_d})$. Let $\pi_i:\Sigma(\mathbf{m})\to$

$\mathbb{P}(\mathbb{C}^{m_i})$ be the projection on the i-th component. Then $\pi_i^* H(m_i), \pi_i^* Q(m_i)$ are the pull back of the bundles $H(m_i), Q(m_i)$ on $\mathbb{P}(\mathbb{C}^{m_i})$ to $\Sigma(\mathbf{m})$ respectively. Denote $t_i = c_1(\pi_i^* H(m_i))$. The cohomology ring $H^*(\Sigma(\mathbf{m}))$ is generated by $t_1, \ldots, t_d$ with the relations $t_i^{m_i} = 0$, that is $H^*(\Sigma(\mathbf{m})) \simeq \mathbb{C}[t_1, \ldots, t_d]/(t_1^{m_1}, \ldots, t_d^{m_d})$ and in the following we interpretate t_i just as variables. Correspondingly, the Chern polynomial of a vector bundle E over $\Sigma(\mathbf{m})$ can be written as $C(t, E) = \tilde{C}(t, \ldots, t, E)$ where $\tilde{C}(t_1, \ldots, t_d, E) = \sum_\alpha c_\alpha t_1^{\alpha_1} \ldots t_d^{\alpha_d}$, $\alpha = (\alpha_1, \ldots, \alpha_d)$ and $c_\alpha \in \mathbf{Z}$. The i-th Chern class $c_i(E)$ coincides with the degree i summand $\sum_{|\alpha|=i} c_\alpha t_1^{\alpha_1} \ldots t_d^{\alpha_d}$, where $|\alpha| = \sum_{i=1}^d \alpha_i$. Since we have $t_i^{m_i} = 0$, note that the top Chern class coincides with $c_{m_1-1,\ldots,m_d-1}$

Let $R(i, \mathbf{m})$ and $R(\mathbf{m})$ be the following vector bundles on $\Sigma(\mathbf{m})$

$$R(i, \mathbf{m}) = (\otimes_{j \in [d] \setminus \{i\}} \pi_j^* H(m_j)) \otimes \pi_i^* Q(m_i), \quad R(\mathbf{m}) = \oplus_{i \in [d]} R(i, \mathbf{m}). \tag{2.2}$$

Note that rank $R(i, \mathbf{m}) = m_i - 1$ and rank $R(\mathbf{m}) = \dim \Sigma(\mathbf{m})$. From the Euler sequence

$$0 \longrightarrow (\otimes_{j \in [d] \setminus \{i\}} \pi_j^* H(m_j)) \otimes \pi_i^* T(m_i) \longrightarrow (\otimes_{j \in [d] \setminus \{i\}} \pi_j^* H(m_j)) \otimes \mathbb{C}^{m_i} \longrightarrow R(i, \mathbf{m}) \longrightarrow 0$$

and from Whitney's formula yield

$$C(t, R(i, \mathbf{m})) = \frac{\left(1 + \sum_{k \in [d] \setminus \{i\}} t_k\right)^{m_i}}{1 - t_i + \sum_{k \in [d] \setminus \{i\}} t_k} \tag{2.3}$$

$$C(t, R(\mathbf{m})) = \prod_{i \in [d]} C(t, R(i, \mathbf{m})). \tag{2.4}$$

In what follows we need to compute the top Chern class of $R(\mathbf{m})$. Since rank $R(\mathbf{m}) = \dim \Sigma(\mathbf{m})$, and $\Sigma(\mathbf{m})$ is a manifold, it follows that the top Chern class of $R(\mathbf{m})$ is of the form $c(\mathbf{m}) \prod_{i \in [d]} t_i^{m_i - 1}$, the coefficient of $t^{\dim \Sigma(\mathbf{m})}$ in $C(t, R(\mathbf{m}))$. By abuse of language the coefficient $c(\mathbf{m})$, which is a positive integer, is called also the top Chern class of $R(\mathbf{m})$.

Lemma 1 *Let $R(\mathbf{m})$ be the vector bundle on the Segre variety $\Sigma(\mathbf{m})$ given by* (2.2)*. Then the top Chern class of $\mathbb{R}(\mathbf{m})$, $c(\mathbf{m})$, is the coefficient of the monomial $\prod_{i \in [d]} t_i^{m_i - 1}$ in the polynomial*

$$\prod_{i \in [d]} \left(\sum_{j=0}^{m_i - 1} \left(\sum_{k \in [d] \setminus \{i\}} t_k \right)^j t_i^{m_i - 1 - j} \right). \tag{2.5}$$

Proof. Call $\hat{t_i} = \sum_{k \in [d] \setminus \{i\}} t_k$.

Write the contribution of the denominator of (2.3) as

$$\frac{1}{1 - t_i + \sum_{k \in [d] \setminus \{i\}} t_k} = \sum_{j \geq 0} (t_i - \hat{t_i})^j$$

expand the numerator of (2.3) as

$$\left(1 + \sum_{k \in [d] \setminus \{i\}} t_k\right)^{m_i} = \sum_{j=0}^{m_i} \binom{m_i}{j} \hat{t_i}^j$$

Putting together, the summand of total degree $m_i - 1$ in (2.3) is

$$\sum_{j=0}^{m_i-1} \binom{m_i}{j+1} \hat{t_i}^{m_i-j-1} (t_i - \hat{t_i})^j = \frac{\sum_{j=0}^{m_i-1} \binom{m_i}{j+1} \hat{t_i}^{m_i-j-1} (t_i - \hat{t_i})^{j+1}}{t_i - \hat{t_i}} = \frac{\left[\hat{t_i} + (t_i - \hat{t_i})\right]^{m_i} - \hat{t_i}^{m_i}}{t_i - \hat{t_i}}$$

which is equal to

$$\sum_{j=0}^{m_i-1} (\hat{t_i})^j t_i^{m_i-1-j}$$

In conclusion, the top Chern class of $R(i, \mathbf{m})$ is given by the coefficient of total degree $m_i - 1$ in the polynomial $\sum_{j=0}^{m_i-1} (\sum_{k \in [d] \setminus \{i\}} t_k)^j t_i^{m_i-1-j} (t_j - \hat{t_j})^i$. The identity (2.4) is equivalent to the claim. $\square$

Note that the polynomial (2.5) is equal to the polynomial given in (1.3).

2.5 Bertini's type theorem

Let M be a compact complex manifold and E a holomorphic bundle on M. A holomorphic section σ of E on open $U \subset E$ is a holomorphic map $\sigma : U \to E$, where E is viewed as a complex manifold. Specifically, let $U_i, i \in [N]$ be the finite cover of M such that the bundle E restricted to U_i is $U_i \times \mathbb{C}^k$ with the standard basis $[\mathbf{e}_{1,i}, \ldots, \mathbf{e}_{k,i}]$, as in §2.2. Then $\sigma(\zeta) = \sum_{j=1}^{k} \sigma_{j,i}(\zeta) \mathbf{e}_{j,i}$ for $\zeta \in U \cap U_i$. where $\sigma_{j,i}(\zeta), j \in [k]$ are analytic on $U \cap U_i$. σ is called a global section if $U = M$. Denote by $\mathrm{H}^0(E)$ the linear space of global sections on E. A subspace $\mathbf{V} \subset \mathrm{H}^0(E)$ is said to generate E if $\mathbf{V}(\zeta)$, the values of all section in $\mathbf{V}$ at each $\zeta \in M$ is equal to E_ζ.

The following proposition is a generalization of the classical Bertini's theorem in algebraic geometry, and it is a standard consequence of the Generic Smoothness Theorem. For the convenience of the reader we state and give a short proof of this proposition.

Theorem 2 ("Bertini's type" theorem) *Let E be a vector bundle on M. Let $\mathbf{V} \subset \mathrm{H}^0(E)$ be a subspace which generates E. Then*

- *if* $\operatorname{rank} E > \dim M$ *for the general* $\sigma \in \mathbf{V}$ *the zero locus of* σ *is empty.*
- *if* $\operatorname{rank} E \leq \dim M$ *for the general* $\sigma \in \mathbf{V}$ *the zero locus of* σ *is either smooth of codimension* $\operatorname{rank} E$ *or it is empty.*
- *if* $\operatorname{rank} E = \dim M = n$ *the zero locus of the general* $\sigma \in V$ *consists of* $c_n(E)$ *simple points, where* c_n *is the* n*-th Chern class, (the top Chern class) and we identify* $H^{2n}(M) = \mathbf{Z}$.

Proof. We identify the vector bundle E with its locally free sheaf of sections, see [6] B.3. We have the projection $E \xrightarrow{\pi} X$, where the fiber $\pi^{-1}(\zeta)$ is isomorphic to the vector space E_ζ. Let $\Sigma \subset E$ be the zero section. By the assumption we have a natural projection of maximal rank

$$M \times \mathbf{V} \xrightarrow{p} E.$$

Let $Z = p^{-1}(\Sigma)$, then Z is isomorphic to the variety $\{(\zeta, \sigma) \in E \times \mathbf{V} | \sigma(\zeta) = 0\}$ and it has dimension equal to $\dim M + \dim \mathbf{V} - \operatorname{rank} E$. Consider the natural projection

$Z\xrightarrow{q}\mathbf{V}$, now $\forall\sigma\in V$ the fiber $q^{-1}(\sigma)$ is naturally isomorphic to the zero locus of σ. We have two cases. If q is dominant (namely the image of q is dense) then by the Generic Smoothness Theorem ([10] Corol. III 10.7) $q^{-1}(\sigma)$ is smooth of dimension $\dim X - \operatorname{rank} E$ for general σ.

If q is not dominant (and this always happens in the case $\operatorname{rank} E > \dim M$) then $q^{-1}(\sigma)$ is empty for general σ. This concludes the proof of the first two parts. The third part follows by [6] example 3.2.16. $\square$

3 The number of singular vector tuples of a generic tensor

In this section we compute the number of singular vector tuples of a general tensor $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$. As pointed in §1 a singular value tuple is viewed as a point $([\mathbf{x}_1],\dots,[\mathbf{x}_d])\in\Sigma(\mathbf{m})$. With a slight abuse of notation we identify a vector $\mathbf{x}_j\in\mathbb{C}^{m_j}\setminus\{\mathbf{0}\}$ with the line $[\mathbf{x}_j]$ in $\mathbb{C}^{m_j}$, which is a point in the projective space $\mathbb{P}(\mathbb{C}^{m_j})$.

Lemma 3 *Let $R(i,\mathbf{m}), i\in[d]$ and $R(\mathbf{m})$ be the vector bundles over the Segre variety $\Sigma(\mathbf{m})$ defined in §2.4. Denote by $\mathrm{H}^0(\mathbb{R}(i,\mathbf{m})), i\in[d]$ and $\mathrm{H}^0(R(\mathbf{m}))$ the linear space of global section of $R(i,\mathbf{m}), i\in[d]$ and $R(\mathbf{m})$ respectively. Then the following conditions hold.*

1. *For each $i\in[d]$ there exists a monomorphism $L_i:\mathbb{C}^{\mathbf{m}}\to\mathrm{H}^0(R(i,\mathbf{m}))$ such that $L_i(\mathbb{C}^{\mathbf{m}})$ generates $R(i,\mathbf{m})$ (see §2.5).*
2. *$L=(L_1,\dots,L_d)$ is a monomorphism of the direct sum of d copies of $\mathbb{C}^{\mathbf{m}}$ ($\oplus^d\mathbb{C}^{\mathbf{m}}$) to $\mathrm{H}^0(R(\mathbf{m}))$.*
3. *$\delta:\mathbb{C}^{\mathbf{m}}\to\oplus^d\mathbb{C}^{\mathbf{m}}$ be the diagonal map $\delta(\mathcal{T})=(\mathcal{T},\dots,\mathcal{T})$. Then $L\circ\delta$ is a monomorphism of $\mathbb{C}^{\mathbf{m}}$ to $\mathrm{H}^0(R(\mathbf{m}))$. Furthermore $L\circ\delta(\mathbb{C}^{\mathbf{m}})$ generates $R(\mathbf{m})$.*

Proof. Let $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ and $\mathbf{u}_j\in\mathbb{C}^{m_j}\setminus\{\mathbf{0}\}, i\in[d]$. For $\mathbf{x}\in\mathbb{C}^{m_i}$ let $[[\mathbf{x}]]$ be the corresponding element in the quotient space $\mathbb{C}^{m_i}/[\mathbf{u}_i]$ and consider u_j as a global section of the hyperplane line bundle $H(m_j)$. View $(\times_{j\in[d]\setminus\{i\}}\mathbf{u}_j)\otimes[[\mathbf{x}]]$ as an element in the fiber $R(i,\mathbf{m})_{(\mathbf{u}_1,\dots,\mathbf{u}_d)}$. Then

$$L_i(\mathcal{T})((\mathbf{u}_1,\dots,\mathbf{u}_d)) := (\otimes_{j\in[d]\setminus\{i\}}\mathbf{u}_j)\otimes[[\mathcal{T}\times\otimes_{j\in[d]\setminus\{i\}}]]. \tag{3.1}$$

It is straightforward to check that $L_i(\mathcal{T})$ is a global section of $R(i,\mathbf{m})$.

Assume $\mathcal{T}\neq 0$. Then there exist $\mathbf{v}_j\in\mathbb{C}^{m_j}, j\in[d]$ such that $\mathcal{T}\times\otimes_{j\in[d]}\mathbf{v}_j\neq 0$. Hence $\mathbf{x}:=\mathcal{T}\times\otimes_{j\in[d]\setminus\{i\}}\in\mathbb{C}^{m_i}\setminus\{\mathbf{0}\}$. Let $\mathbf{u}_j=\mathbf{v}_j$ for $j\neq i$. Choose $\mathbf{u}_i\in\mathbb{P}(\mathbb{C}^{m_i})\setminus\{[\mathbf{x}]\}$. Then $L(\otimes_{j\in[d]}\mathbf{v}_j)((\mathbf{u}_1,\dots,\mathbf{u}_d))\neq 0$. Hence L_i is injective.

Let $\mathbf{v}_j=\|\mathbf{u}_j\|^{-1}\mathbf{u}_j^*$ for $j\in[d]\setminus\{i\}$ and $\mathbf{v}_i=\mathbf{x}$. Then

$$L_i(\otimes_{j\in[d]}\mathbf{v}_j)((\mathbf{u}_1,\dots,\mathbf{u}_d)) = (\prod_{j\in[d]\setminus\{i\}}\mathbf{u}_j)\otimes[[\mathbf{x}]].$$

Hence $L(\mathbb{C}^{\mathbf{m}})$ generates $R(i,\mathbf{m})$.

Define $L((\mathcal{T}_1,\dots,\mathcal{T}_d))((\mathbf{u}_1,\dots,\mathbf{u}_d))=\oplus_{i\in[d]}L_i((\mathcal{T}_1,\dots,\mathcal{T}_d))((\mathbf{u}_1,\dots,\mathbf{u}_d))$. Then $L((\mathcal{T}_1,\dots,\mathcal{T}_d))\in\mathrm{H}^0(R(\mathbf{m}))$. Clearly L is a monomorphism.

It is left to show that $L\circ\delta(\mathbb{C}^{\mathbf{m}})$ generates $R(\mathbf{m})$. By changing bases in $\mathbb{C}^{m_1},\ldots,\mathbb{C}^{m_d}$ we may assume with a loss of generality that $\mathbf{u}_i=(1,0,\ldots,0)^\top\in\mathbb{C}^{m_i}$ for $i\in[d]$. Then point each $[[\mathbf{x}_i]]\in\mathbb{C}^{m_i}/[\mathbf{u}_i]$ can be chosen of the form $\mathbf{x}_i=(0,x_{2,i},\ldots,x_{m_i,i})^\top$. Choose $\mathcal{T}=[t_{j_1,\ldots,j_d}]$ as follows. $t_{1,1,\ldots,1}=1$. For a fixed $i\in[d]$ let $t_{1,\ldots,1,j_i,1,\ldots,1}=x_{j_i,i}$ for $j_i\ge 2$. All other entries of $\mathcal{T}$ are zero. Clearly

$$L_i(\mathcal{T})((\mathbf{u}_1,\ldots,\mathbf{u}_d))=(\otimes_{j\in[d]\setminus\{i\}}\mathbf{u}_j)\otimes[[\mathbf{x}_i]],\quad i\in[d].$$

Hence $L\circ\delta(\mathbb{C}^{\mathbf{m}})$ generates $R(\mathbf{m})$. $\square$

It follow from Künneth formula([12]) the equalities

$$L_i(\mathbb{C}^{\mathbf{m}})=\mathrm{H}^0(R(i,\mathbf{m})),\ i\in[d],\quad L(\oplus^d\mathbb{C}^{\mathbf{m}})=H^0(R(\mathbf{m})).\tag{3.2}$$

Assume that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is a a singular vector tuple of $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$, i.e. (1.2) holds. We say that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ corresponds to zero or nonzero singular value if either $\prod_{j\in[d]}\lambda_i$ is zero or nonzero respectively. For $\mathcal{T}\in\mathbb{R}^{\mathbf{m}}$ with a real singular vector tuple $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma_{\mathbb{R}}(\mathbf{m})$ the condition $\prod_{j\in[d]}\lambda_i=0$ implies that $\lambda_i=0$ for each i. Indeed, since $\mathbf{x}_i\in\mathbb{R}^{m_i}\setminus\{\mathbf{0}\}$ it follows from (1.2) that $\lambda_i=\frac{\mathcal{T}\times\otimes_{j\in[d]}\mathbf{x}_j}{\mathbf{x}_i^\top\mathbf{x}_i}$ for each $i\in[d]$. Thus $\lambda_k=0$ for some $k\in[d]$ yields that $\mathcal{T}\times\otimes_{j\in[d]}\mathbf{x}_j=0$. Hence each $\lambda_i=0$.

However, this observation is not valid for complex tensors, already in the case of complex valued matrices ($d=2$), see an example below. It is straightforward to see that a singular value pair $(\mathbf{x}_1,\mathbf{x}_2)$ of $A\in\mathbb{C}^{m_1\times m_2}$ is given by the following conditions

$$A\mathbf{x}_2=\lambda_1\mathbf{x}_1,\quad A^\top\mathbf{x}_1=\lambda_2\mathbf{x}_2,\quad \mathbf{x}_i\in\mathbb{C}^{m_i}\setminus\{\mathbf{0}\},\lambda_i\in\mathbb{C},\quad i=1,2.\tag{3.3}$$

Consider the following simple example.

$$A=\begin{bmatrix}1 & \mathbf{i}\\ -\mathbf{i} & 1\end{bmatrix},\ \mathbf{x}_1=\begin{bmatrix}1\\0\end{bmatrix},\ \mathbf{x}_2=\begin{bmatrix}1\\ \mathbf{i}\end{bmatrix}.$$

Then $A^\top\mathbf{x}_1=\mathbf{x}_2, A\mathbf{x}_2=\mathbf{0}$, i.e. $\lambda_1=1,\lambda_2=0$.

A singular vector tuple $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is called a simple singular tuple of $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$, (simple), if there exist open neighborhoods $O_1\subset\Sigma(\mathbf{m}),O_2\subset\mathbb{C}^{\mathbf{m}}$ of $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ and $\mathcal{T}$ respectively, such that each $\mathcal{S}\in O_2$ has exactly one singular vector tuple in O_1.

Theorem 4 *A general $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ has exactly $c(\mathbf{m})$, the coefficient of the monomial $\prod_{i=1}^d t_i^{m_i-1}$ in the polynomial (1.3), simple singular vector tuples which correspond to nonzero singular values. A general $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ does not have a zero singular value. In particular, a general real valued tensor $\mathcal{T}\in\mathbb{R}^{\mathbf{m}}$ has at most $c(\mathbf{m})$ real singular pairs, which are simple.*

Proof. Let $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ and consider the section $\hat{\mathcal{T}}:=L\circ\delta(\mathcal{T})\in\mathrm{H}^0(R(\mathbf{m}))$. We claim that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is a zero of $\hat{\mathcal{T}}$ if and only if $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ is a singular vector tuple corresponding to $\mathcal{T}$. Furthermore, $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is a simple zero of $\hat{\mathcal{T}}$ if and only if $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ is a simple singular vector tuple corresponding to $\mathcal{T}$. Indeed, suppose first that $\hat{T}((\mathbf{x}_1,\ldots,\mathbf{x}_d))=0$. Then $L_i(\mathcal{T})((\mathbf{x}_1,\ldots,\mathbf{x}_d))$ is a zero vector in the fiber $R(i,\mathbf{m})_{(\mathbf{x}_1,\ldots,\mathbf{x}_d)}$. Suppose first that $\mathcal{T}\times(\prod_{j\in[d]\setminus\{i\}}\mathbf{x}_j)\neq\mathbf{0}$. Then $\mathcal{T}\times(\prod_{j\in[d]\setminus\{i\}}\mathbf{x}_j)=\lambda_i\mathbf{x}_i$ for some $\lambda_i\neq 0$. Otherwise the previous equality hods with $\lambda_i=0$. Hence $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ is a singular vector pair corresponding to $\mathcal{T}$. Viceversa, it is straightforward to see that if $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ is a singular vector pair corresponding to $\mathcal{T}$ then the section $\mathcal{T}$ vanishes at $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$.

Since $L\circ\delta(\mathbb{C}^{\mathbf{m}})$ is a generating subspace of $\mathrm{H}^0(R(\mathbf{m}))$ and $\operatorname{rank} R(\mathbf{m})=\dim\Sigma(\mathbf{m})$, Bertini's type theorem, see §2.5, yields that for a general $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ the section $L\circ\delta(\mathcal{T})$ has a finite number of simple zeros, and this number is equal to the top Chern class of $R(\mathbf{m})$. Lemma 3 yields that the top Chern class of $R(\mathbf{m})$ is $c(\mathbf{m})$.

It is straightforward to show that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is a simple zero $\hat{\mathcal{T}}$ if and only if it is an isolated zero and there exist open neighborhoods $O_1\subset\Sigma(\mathbf{m}),O_2\subset\mathbb{C}^{\mathbf{m}}$ of $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ and $\mathcal{T}$ respectively, such that for each $\mathcal{S}\in O_2$ the section $\hat{\mathcal{S}}$ has exactly one singular vector tuple in O_1. Clearly this fact is equivalent to the assumption that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ is a simple singular vector tuple of $\mathcal{T}$.

It is left to show that a generic $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ does not have a zero singular value. Fix $i\in[d]$ and consider the set of all $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ which have a singular vector tuple $(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\Sigma(\mathbf{m})$ with $\lambda_i=0$. Let $T_i:=\mathbb{C}^{m_i}$ be the trivial vector bundle of rank m_i over $\mathbb{P}(\mathbb{C}^{m_i})$. Denote by $T(i,\mathbf{m}):=\pi_i^*T_i$ the pull back of this trivial bundle to $\Sigma(\mathbf{m})$. Let

$$R(i,\mathbf{m})':=(\otimes_{j\in[d]\setminus\{i\}}\pi_j^*H(m_j))\otimes T(i,\mathbf{m}),\ R_i(\mathbf{m})':=(\oplus_{j\in[d]\setminus\{i\}}R(j,\mathbf{m}))\oplus R(i,\mathbf{m})'.$$

Note that $R_i(\mathbf{m})'$ is a vector bundle on $\Sigma(\mathbf{m})$ of rank $\dim\Sigma(\mathbf{m})+1$. As above we can define a monomorphism $L_i':\mathbb{C}^{\mathbf{m}}\to\mathrm{H}^0(R(i,m)')$ defined by the equality

$$L_i'(\mathcal{T})((\mathbf{u}_1,\ldots,\mathbf{u}_d)):=(\otimes_{j\in[d]\setminus\{i\}}\mathbf{u}_j)\otimes(\mathcal{T}\times\otimes_{j\in[d]\setminus\{i\}}\mathbf{u}_j).$$

Let $\tilde{L}_i=(L_1,\ldots,L_{i-1},L_i',L_{i+1},\ldots,L_d):\oplus_{j\in[d]}\mathbb{C}^{\mathbf{m}}\to\mathrm{H}^0(R_i(\mathbf{m})')$. We claim that $\tilde{L}_i\circ\delta(\mathbb{C}^{\mathbf{m}})$ generates $R_i(\mathbf{m})$. It is enough to consider the point $(\mathbf{u}_1,\ldots,\mathbf{u}_d)\in\Sigma(\mathbf{m})$, where $\mathbf{u}_i=(1,0,\ldots,0)^\top\in\mathbb{C}^{m_i}$ for $i\in[d]$. Let $\mathbf{x}_j=(x_{1,j},\ldots,x_{m_j,j})^\top\in\mathbb{C}^{m_j}$ for $j\in[d]$. Choose $\mathcal{T}=[t_{j_1,\ldots,j_d}]$ as follows. $t_{1,1,\ldots,1}=x_{1,i}$. For a fixed $j\in[d]$ let $t_{1,\ldots,1,k_j,1,\ldots,1}=x_{k_j,j}$ for $k_j\geq 2$. All other entries of $\mathcal{T}$ are zero. Then

$$\begin{aligned}&L_j(\mathcal{T})((\mathbf{u}_1,\ldots,\mathbf{u}_d))=(\otimes_{k\in[d]\setminus\{j\}}\mathbf{u}_k)\otimes[[\mathbf{x}_j]],\quad j\in[d]\setminus\{i\},\\&\tilde{L}_i(\mathcal{T})((\mathbf{u}_1,\ldots,\mathbf{u}_d))=(\otimes_{k\in[d]\setminus\{i\}}\mathbf{u}_k)\otimes\mathbf{x}_i.\end{aligned}$$

Hence $\tilde{L}_i\circ\delta(\mathbb{C}^{\mathbf{m}})$ generates $R_i(\mathbf{m})'$. Theorem 2 yields that for general $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ the section $\tilde{L}_i\circ\delta(\mathcal{T})$ has no zeros. That is, $\mathcal{T}$ does not have a singular vector tuple satisfying (1.2) with $\lambda_i=0$. Hence a general tensor $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}$ does not have a zero singular value.

In algebraic geometry the term general $\mathcal{T}$ means that $\mathcal{T}\in\mathbb{C}^{\mathbf{m}}\setminus X$, where X is some algebraic variety, given as a zero set of a finite number of polynomial in the entries of $\mathcal{T}$. A general real $\mathcal{T}\in\mathbb{R}^{\mathbf{m}}$ does not lie in X. Hence a general $\mathcal{T}\in\mathbb{R}^{\mathbf{m}}$ has exactly $c(\mathbf{m})$ simple singular value tuples. Only some of those can be realized as points in $\Sigma_{\mathbb{R}}(\mathbf{m})$. □

We first observe that Theorem 4 agrees with the standard theory of singular values for $m\times n$ real matrices. Let $A\in\mathbb{C}^{m_1\times m_2}$ and $U_i\in\mathbb{C}^{m_i\times m_i}$ (for $i=1,2$) be (complex) orthogonal matrices. Suppose that (3.3) holds. Let $B=U_1AU_2^\top$. Then

$(U_1\mathbf{x}_1, U_2\mathbf{x}_2)$ is a singular pair of B. Suppose that there exists U_1, U_2 such that B is a diagonal matrix such that the absolute values of its diagonal entries are positive and pairwise distinct. Then a straighforward calculation shows that B has exactly $\min(m_1, m_2)$ singular pairs corresponding to nonzero singular values. The singular value decomposition of $A \in \mathbb{R}^{m_1 \times m_2}$ implies that a general $A \in \mathbb{R}^{m_1 \times m_2}$ has exactly $\min(m_1, m_2)$ singular pairs corresponding to nonzero singular values. It is also easy to see that $c(m_1, m_2) = \min(m_1, m_2)$.

We point out a matrix proof of Theorem 4 for $d = 2$. Let $\mathrm{O}(m) \subset \mathbb{C}^{m \times m}$ be variety of $m \times m$ orthogonal matrices and $\mathrm{D}_{m,n} \subset \mathbb{C}^{m \times n}$ the linear subspace of all diagonal matrices. Consider the trilinear polynomial map $(U_1, D, U_2) \mapsto U_1^\top D U_2$ from the $\mathrm{O}(m_1) \times \mathrm{D}_{m_1,m_2} \times \mathrm{O}(m_2)$ to $\mathbb{C}^{m_1 \times m_2}$. It is not difficult to show that the image of this map is dense in $\mathbb{C}^{m_1 \times m_2}$, since over the reals this map is onto. The above arguments give a simple proof of Theorem 4 for $d = 2$.

We list for the convenience of the reader a few values $c(\mathbf{m})$. First

$$c(\underbrace{2, \ldots, 2}_{d}) = d! \tag{3.4}$$

Second, we list at next page the first values in the case $d = 3$. ¿From this table one sees that $c(m_1, m_2, m_3)$ stabilizes for $m_3 \geq m_1 + m_2 - 1$, the case when equality holds is called the boundary format case in the theory of hyperdeterminants ([7]). It is the case where a “diagonal” naturally occurs, like in the following figure



In $d = 2$ case, boundary format means square.

d_1, d_2, d_3	$c(d_1, d_2, d_3)$	
$2, 2, 2$	6	
$2, 2, n$	8	$n \geq 3$
$2, 3, 3$	15	
$2, 3, 4$	18	$n \geq 4$
$2, 4, 4$	28	
$2, 4, n$	32	$n \geq 5$
$2, 5, 5$	45	
$2, 5, n$	50	$n \geq 6$
$2, m, m+1$	$2m^2$	
$3, 3, 3$	37	
$3, 3, 4$	55	
$3, 3, n$	61	$n \geq 5$
$3, 4, 4$	104	
$3, 4, 5$	138	
$3, 4, n$	148	$n \geq 6$
$3, 5, 5$	225	
$3, 5, 6$	280	
$3, 5, n$	295	$n \geq 7$
$3, m, m+2$	$\frac{8}{3}m^3 - 2m^2 + \frac{7}{3}m$	
$4, 4, 4$	240	
$4, 4, 5$	380	
$4, 4, 6$	460	
$4, 4, n$	480	$n \geq 7$
$4, 5, 5$	725	
$4, 5, 6$	1030	
$4, 5, 7$	1185	
$4, 5, n$	1220	$n \geq 8$
$5, 5, 5$	1621	
$5, 5, 6$	2671	
$5, 5, 7$	3461	
$5, 5, 8$	3811	
$5, 5, n$	3881	$n \geq 9$

4 Partially symmetric singular value tuples

For an integer $m \geq 2$ let $m^{\times d} := (\underbrace{m, \ldots, m}_{d})$. Then $\mathcal{T} \in \mathbb{F}^{m^{\times d}}$ is called d-cube, or simply a cube tensor. Denote by $\mathrm{S}^d(\mathbb{F}^m) \subset \mathbb{F}^{m^{\times d}}$ the subspace of symmetric tensors. For $\mathcal{T} \in \mathrm{S}^d(\mathbb{F}^m)$ it is natural to consider a singular tuple (1.2) where $\mathbf{x}_1 = \ldots = \mathbf{x}_d = \mathbf{x}$ [14, formula (7) with $p = 2$]. This is equivalent to the system

$$\mathcal{T} \times \otimes^{d-1}\mathbf{x} = \lambda \mathbf{x}, \quad \mathbf{x} \neq 0. \tag{4.1}$$

Here $\otimes^{d-1}\mathbf{x} := \underbrace{\mathbf{x} \otimes \ldots \otimes \mathbf{x}}_{d-1}$. Furthermore, the contraction in (4.1) is on the last $d-1$ indices. The equation (4.1) makes sense for any cube tensor $\mathcal{T} \in \mathbb{C}^{m^{\times d}}$ [14, 17, 20]. For $d = 2$ $\mathbf{x}$ is an eigenvector of the square matrix $\mathcal{T}$. Hence for a d-cube tensor

($d \geq 3$) $\mathbf{x}$ is referred as a nonlinear eigenvalue of $\mathcal{T}$. Abusing slightly our notation we call $(\mathbf{x}, \ldots, \mathbf{x}) \in \Sigma(m^{\times d})$ a symmetric singular tuple of $\mathcal{T}$. (Note that if $\mathcal{T} \in \mathrm{S}^d(\mathbb{C}^m)$ then $(\mathbf{x}, \ldots, \mathbf{x})$ is a proper symmetric singular tuple of $\mathcal{T}$.)

Let $s_{d-1}(\mathcal{T}) = [t'_{i_1,\ldots,i_d}]$ be the symmetrization of a d-cube $\mathcal{T} = [t_{i_1,\ldots,i_d}]$ with respect to the last $d-1$ indices

$$t'_{i_1,\ldots,i_d} = \frac{1}{p(i_2,\ldots,i_d)} \sum_{\{j_2,\ldots,j_d\}=\{i_2,\ldots,i_d\}} t_{i_1,j_2,\ldots,j_d}. \tag{4.2}$$

Here $p(i_2,\ldots,i_d)$ is the number of multisets $\{j_2,\ldots,j_d\}$ which are equal to $\{i_2,\ldots,i_d\}$. (Note that for $d=2$ $s_1(\mathcal{T}) = \mathcal{T}$.) It is straightforward to see that

$$\mathcal{T} \times \otimes^{d-1}\mathbf{y} = s_{d-1}(\mathcal{T}) \otimes^{d-1} \mathbf{y} \text{ for all } \mathbf{y}. \tag{4.3}$$

Hence in (4.1) we can assume that $\mathcal{T}$ is symmetric with respect to the last $d-1$ indices.

As for singular value tuples we view the eigenvectors of $\mathcal{T}$ as elements of $\mathbb{P}(\mathbb{C}^m)$. It was shown by Cartwright and Sturmfels [2] that a general $\mathcal{T} \in \mathbb{C}^{m^{\times d}}$ has exactly $\frac{(d-1)^m-1}{d-2}$ distinct eigenvectors. (This formula was conjectured in [17].)

The aim of this section is to consider "partially symmetric singular vectors" and their numbers for a general tensor. This number will interpolate our formula $c(\mathbf{m})$ for the number of singular tuples for a general $\mathcal{T} \in \mathbb{C}^{\mathbf{m}}$ and the number of eigenvalues of general $\mathcal{T} \in \mathbb{C}^{m^{\times d}}$ given in [2].

Let $d = \omega_1 + \ldots + \omega_p$ be a partition of d. So each ω_i is a positive integer. Let $\omega_0 = m'_0 = 0$, $\boldsymbol{\omega} = (\omega_1, \ldots, \omega_p)$ and denote by $\mathbf{m}(\boldsymbol{\omega})$ the d-tuple

$$\mathbf{m}(\boldsymbol{\omega}) = (\underbrace{m'_1,\ldots,m'_1}_{\omega_1},\ldots,\underbrace{m'_p,\ldots,m'_p}_{\omega_p}) = (m_1,\ldots,m_d). \tag{4.4}$$

Denote by $\mathrm{S}^{\boldsymbol{\omega}}(\mathbb{F}) \subset \mathbb{F}^{\mathbf{m}(\boldsymbol{\omega})}$ the subspace of tensors which are partially symmetric with respect to the partition $\boldsymbol{\omega}$. That is the entries of $\mathcal{T} = [t_{i_1,\ldots,i_d}] \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{F})$ are invariant, if we permute indices in the $k-th$ group of indices $[\sum_{j=0}^{k} \omega_j] \setminus [\sum_{j=0}^{k-1} \omega_j]$ for $k \in [p]$. Note that $\mathrm{S}^{\boldsymbol{\omega}}(\mathbb{F}) = \mathrm{S}^d(\mathbb{F}^{\mathbf{m}}))$ for $p=1$ and $\mathrm{S}^{\boldsymbol{\omega}}(\mathbb{F}) = \mathbb{F}^{\mathbf{m}}$ for $p=d$. We call $\boldsymbol{\omega} = (1,\ldots,1)$, i.e. $p=d$, the *trivial* partition.

For simplicity of notation we let $\mathrm{S}^{\boldsymbol{\omega}} := \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{C})$. Assume that $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$. Consider a singular vector tuple $(\mathbf{x}_1, \ldots, \mathbf{x}_d)$ satisfying (1.2) and $\boldsymbol{\omega}$-symmetric conditions

$$\mathbf{x}_j = \mathbf{y}_k \text{ for } j \in [\sum_{i=1}^{k} \omega_i m'_i] \setminus [\sum_{i=0}^{k-1} \omega_i m'_i],\ k \in [p]. \tag{4.5}$$

We rewrite (1.2) for an $\boldsymbol{\omega}$-symmetric singular vector tuple $(\mathbf{x}_1, \ldots, \mathbf{x}_d)$ as follows. Define

$$\otimes_{l\in[p]} (\otimes^{\omega_l-\delta_{lk}} \mathbf{y}_l) := \otimes_{j\in[d]\setminus\{1+\sum_{i=0}^{k-1}\omega_i m'_i\}} \mathbf{x}_j, \text{ for } k \in [p]. \tag{4.6}$$

Hence our equations for an $\boldsymbol{\omega}$ symmetric singular value tuple for $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$ is given by

$$\mathcal{T} \times \otimes_{l\in[p]}(\otimes^{\omega_l-\delta_{lk}} \mathbf{y}_l) = \lambda_k \mathbf{y}_k \quad k \in [p]. \tag{4.7}$$

In view of the definition of $\otimes_{l\in[p]}(\otimes^{\omega_l-\delta_{lk}}\mathbf{y}_l)$ we agree that the contraction on the left-hand side of (4.7) is done on all indices except the index $1+\sum_{i=0}^{k-1}\omega_i m_i'$. As for the d-cube tensor the system (4.7) makes sense for any $\mathcal{T}\in\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}$.

Let $\mathbf{m}' := (m_1',\ldots,m_p')$. We call $(\mathbf{y}_1,\ldots,\mathbf{y}_p)\in\Sigma(\mathbf{m}')$ satisfying (4.7) as $\boldsymbol{\omega}$-symmetric singular tuples of $\mathcal{T}\in\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}$. We say that $(\mathbf{y}_1,\ldots,\mathbf{y}_p)$ corresponds to a zero (nonzero) singular value if $\prod_{i=1}^{p}\lambda_i = 0$ $(\neq 0)$.

The aim of this section to show that a general tensor $\mathcal{T}\in\mathrm{S}^{\boldsymbol{\omega}}$, hence a general tensor $\mathcal{T}\in\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}$, has exactly $c(\mathbf{m}',\boldsymbol{\omega})$ $\boldsymbol{\omega}$-symmetric singular tuples, where $c(\mathbf{m}',\boldsymbol{\omega})$ is the coefficient of the monomial $\prod_{i\in[p]}t_i^{m_i'-1}$ in the homogeneous polynomial

$$\prod_{i\in[p]}\frac{\hat{t}_i^{m_i'}-t_i^{m_i'}}{\hat{t}_i-t_i},\quad \hat{t}_i=(\omega_i-1)t_i+\sum_{j\in[p]\setminus\{i\}}\omega_j t_j,\ i\in[p]. \tag{4.8}$$

The proof of this result is analogous to the proof of Theorem 4, so we point briefly the needed modifications. Let $H(m_i')$ and $Q(m_i')$ be the vector bundles defined in §2.4. Let π_i be the projection of $\Sigma(\mathbf{m}')$ on the component $\mathbb{P}(\mathbb{C}^{m_i'})$. Then $\pi_i^*H(m_i'),\pi_i^*Q(m_i')$ are the pull backs of the vector bundles $H(m_i'),Q(m_i')$ to $\Sigma(\mathbf{m}')$ respectively. Clearly $C(t,\pi_i^*H(m_i'))=1+t_i$ and moreover $C(t,\otimes^k\pi_i^*H(m_i'))=1+kt_i$, where $t_i^{m_i'}=0$. Define the following vector bundles on $\Sigma(\mathbf{m}')$.

$$\begin{aligned}&\tilde{R}(i,\mathbf{m}') := \Big(\otimes_{j\in[p]}(\otimes^{\omega_j-\delta_{ji}}\pi_j^*H(m_j'))\Big)\otimes\pi_i^*Q(m_i'),\ i\in[p],\\ &\tilde{R}(\mathbf{m}') := \oplus_{i\in[p]}\tilde{R}(i,\mathbf{m}').\end{aligned} \tag{4.9}$$

Note that rank $\tilde{R}(i,\mathbf{m}')=m_i'-1$ and rank $\tilde{R}(\mathbf{m}')=\dim\Sigma(\mathbf{m}')$. As in the proof of Lemma 1 we deduce that the top Chern class of $\tilde{R}(i,\mathbf{m}')$ is given by the polynomial

$$\sum_{j=0}^{m_i'-1}\Big(\sum_{k\in[p]}(\omega_k-\delta_{ki})t_k\Big)^j t_i^{m_i'-1-j}),\quad i\in[p], \tag{4.10}$$

where we assume the relations $t_i^{m_i'}=0$ for $i\in[p]$. Use Whitney's formula to deduce that the top Chern class of $\tilde{R}(\mathbf{m}')$ is $c(\mathbf{m}',\boldsymbol{\omega})$.

We next observe that we can view $\Sigma(\mathbf{m}')$ as a submanifold by using the imbedding $\eta:\Sigma(\mathbf{m}')\to\Sigma(\mathbf{m}(\boldsymbol{\omega}))$. Here $\eta((\mathbf{y}_1,\ldots,\mathbf{y}_p))=(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ where we assume the relations (4.5). It now follows that $\tilde{R}(i,\mathbf{m}')$ is a pull back $R(j,\mathbf{m})$, where $j=1+\sum_{k=0}^{i-1}\omega_k m_k'$. From the results of §3, in particular Lemma 3, we deduce that there exists a monomorphism $L_i:\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}\to\mathrm{H}^0(\tilde{R}(i,\mathbf{m}'))$. Furthermore $L_i(\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})})$ generates $\tilde{R}(i,\mathbf{m}')$.

Let $L=(L_1,\ldots,L_p):\oplus_{i\in[p]}\mathbb{C}^{\mathbf{m}(\omega)}\to\mathrm{H}^0(\tilde{R}(\mathbf{m}'))$. Then $L\circ\delta(\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})})$ generates $\mathrm{H}^0(\tilde{R}(\mathbf{m}'))$.

Theorem 5 *Let $d\geq 3$ be a an integer and assume that $\boldsymbol{\omega}=(\omega_1,\ldots,\omega_p)$ is a partition of d. Let $\mathbf{m}(\boldsymbol{\omega})$ be defined by (4.6). Denote by $\mathrm{S}^{\boldsymbol{\omega}}\subset\mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}$ the subspace of tensors partially symmetric with respect to $\boldsymbol{\omega}$. A general $\mathcal{T}\in\mathrm{S}^{\boldsymbol{\omega}}$ has exactly $c(\mathbf{m}',\boldsymbol{\omega})$, the coefficient of the monomial $\prod_{i=1}^{p}t_i^{m_i'-1}$ in the polynomial (4.8), simple $\boldsymbol{\omega}$-symmetric singular vector tuples which correspond to nonzero singular values. A general $\mathcal{T}\in\mathrm{S}^{\boldsymbol{\omega}}$ does not have a zero singular value. In particular, a general real valued tensor $\mathcal{T}\in\mathrm{S}_{\mathbb{R}}^{\boldsymbol{\omega}}$ has at most $c(\mathbf{m}',\boldsymbol{\omega})$ real singular pairs, which are simple.*

Proof. The observations given before the statement of the theorem yield all the results for a general $\mathcal{T} \in \mathbb{C}^{\mathbf{m}(\boldsymbol{\omega})}$. Observe next that $L \circ \delta(\mathrm{S}^{\boldsymbol{\omega}})$ generates $\tilde{R}(\mathbf{m}')$. It is enough to show that $L \circ \delta(\mathrm{S}^{\boldsymbol{\omega}})$ generates the fiber $\tilde{R}(\mathbf{m}')$ over the point $(\mathbf{u}_1, \ldots, \mathbf{u}_p)$, where $\mathbf{u}_i = (1, 0, \ldots, 0)^\top \in \mathbb{C}^{m'_i}$ for $i \in [p]$. Each $[[\mathbf{x}_i]] \in \mathbb{C}^{m'_i}/[\mathbf{u}_i]$ can be chosen of the form $\mathbf{x}_i = (0, x_{2,i}, \ldots, x_{m'_i,i})^\top$. Choose $\mathcal{T} = [t_{j_1,\ldots,j_d}]$ as follows. $t_{1,1,\ldots,1} = 1$. For a fixed $i \in [d]$ let $t_{1,\ldots,1,j_{q(i)},1,\ldots,1} = x_{j_q(i),i}$ for $j_{q(i)} \geq 2$, where $q(i) = 1 + \sum_{k=0}^{i-1} m'_k$. Clearly

$$L_i(\mathcal{T})((\mathbf{u}_1, \ldots, \mathbf{u}_p)) = (\otimes_{j\in[p]}(\otimes^{\omega_j - \delta_{ij}} \mathbf{u}_j) \otimes [[\mathbf{x}_i]], \quad i \in [p].$$

Hence $L \circ \delta(S^{\boldsymbol{\omega}})$ generates $\tilde{R}(\mathbf{m}')$. Bertini's theorem yields that a general $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$ has exactly $c(\mathbf{m}', \boldsymbol{\omega})$ simple $\boldsymbol{\omega}$-symmetric singular vector tuples. The proof that a general $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$ does not have a zero singular value is analogous to the proof given in Theorem 4. $\square$

Remark 6 *In the special case* $\boldsymbol{\omega} = (1, 1, \ldots, 1)$ *we have* $c(\mathbf{m}', \boldsymbol{\omega})) = c(\mathbf{m}')$ *and Theorem 5 reduces to Theorem 4. In the case* $\boldsymbol{\omega} = (d)$ *we have* $c(m, \boldsymbol{\omega}) = \frac{(d-1)^m - 1}{d-2}$ *and Theorem 5 reduces to the results in [2]. This last reduction was performed already in [18].*

Lemma 7 *In the case* $\boldsymbol{\omega} = (d-1, 1)$ *we have*

$$c((m_1, m_2), (d-1, 1)) = \sum_{i=0}^{m_1-1} \sum_{j=0}^{m_2-1} \binom{i}{j} (d-2)^j (d-1)^{i-j}.$$

If $m_1 \leq m_2$ *we have* $c((m_1, m_2), (d-1, 1)) = \frac{(2d-3)^{m_1} - 1}{2d-4}$
If $m_1 = m_2 + 1$ *we have* $c((m_1, m_2), (d-1, 1)) = \frac{(2d-3)^{m_1} - 1}{2d-4} - (d-1)^{m_1 - 1}$

Note that in the $3 \times 3 \times 3$ case our formula gives $c(3, 3, 3) = c((3, 3, 3), (1, 1, 1)) = 37$, while Cartwright-Sturmfels formula gives 7, counting indeed triples $(\mathbf{x}, \mathbf{x}, \mathbf{x})$. Note also that the number 37 was computed, in a similar setting, in [16].

Moreover the previous lemma gives $c((3, 3), (2, 1)) = 13$ for the singular vectors of the form $(\mathbf{x}, \mathbf{x}, \mathbf{y})$.

It is instructive to list all the singular vectors for the diagonal tensor $\mathcal{T} = [\delta_{i_1,i_2}\delta_{i_2 i_3}] \in \mathbb{C}^{3\times3\times3}$.

$(x0, x1, x2)(y0, y1, y2)(z0, z1, z2)$	singular value	
$(1, 0, 0)(1, 0, 0)(1, 0, 0)$	1	
$(0, 1, 0)(0, 1, 0)(0, 1, 0)$	1	
$(0, 0, 1)(0, 0, 1)(0, 0, 1)$	1	
$(1, 1, 0)(1, 1, 0)(1, 1, 0)$	1	
$(1, 0, 1)(1, 0, 1)(1, 0, 1)$	1	
$(0, 1, 1)(0, 1, 1)(0, 1, 1)$	1	
$(1, 1, 1)(1, 1, 1)(1, 1, 1)$	1	
$(1, 1, 0)(1, -1, 0)(1, -1, 0)$	1	3 permutations
$(1, 0, 1)(1, 0, -1)(1, 0, -1)$	1	3 permutations
$(0, 1, 1)(0, 1, -1)(0, 1, -1)$	1	3 permutations
$(1, 1, 1)(1, 1 - 1)(1, 1, -1)$	1	3 permutations
$(1, 1, 1)(1, -1, 1)(1, -1, 1)$	1	3 permutations
$(1, 1, 1)(-1, 1, 1)(-1, 1, 1)$	1	3 permutations
$(1, 0, 0)(0, 1, 0)(0, 0, 1)$	0	6 permutations
$(1, 1, -1)(1, -1, 1)(-1, 1, 1)$	-1	6 permutations

In this list the first 7 singular vectors have equal entries and they are the one counted by the formula in [2].

The first $7+6 = 13$ singular vectors have the form $(\mathbf{x}, \mathbf{x}, \mathbf{y})$. Any singular vector of this form gives 3 singular vectors $(\mathbf{x}, \mathbf{x}, \mathbf{y})$, $(\mathbf{x}, \mathbf{y}, \mathbf{x})$, $(\mathbf{y}, \mathbf{x}, \mathbf{x})$.

Note that six singular vectors have zero singular value, but this does not correspond to the general case, indeed for a general tensor all 37 singular vectors correspond to nonzero singular value.

In the case of $4 \times 4 \times 4$ tensors, the diagonal tensor has 156 singular vectors corresponding to nonzero singular value and infinitely many singular vectors corresponding to zero singular vectors. These infinitely many singular vectors fill exactly 36 projective lines in the Segre variety $\mathbb{P}(\mathbb{C}^4) \times \mathbb{P}(\mathbb{C}^4) \times \mathbb{P}(\mathbb{C}^4)$, which "count" in this case for the remaining $240 - 156 = 84$ singular vectors.

5 A homogeneous pencil eigenvalue problem

For $\mathbf{x} = (x_1, \dots, x_m)^\top \in \mathbb{C}^m$ denote $\mathbf{x}^{\circ(d-1)} := (x_1^{d-1}, \dots, x_m^{d-1})^\top$. Let $\mathcal{T} \in \mathbb{C}^{m^{\times d}}$. The eigenvalues of $\mathcal{T}$ satisfying (4.1) are called the E-eigenvalues in [20]. The homogeneous eigenvalue problem introduced in [15], [14] and [19], sometimes referred as N-eigenvalues, is

$$\mathcal{T} \times \otimes^{d-1}\mathbf{x} = \lambda \mathbf{x}^{\circ(d-1)}, \quad \mathbf{x} \neq \mathbf{0}. \tag{5.1}$$

Let $\mathcal{S} \in \mathbb{C}^{m^{\times d}}$. Then a generalized $d-1$ *pencil* eigenvalue problem is

$$\mathcal{T} \times \otimes^{d-1}\mathbf{x} = \lambda \mathcal{S} \times \otimes^{d-1}\mathbf{x}. \tag{5.2}$$

For $d = 2$ the above homogeneous system is the standard eigenvalue problem for a pencil of matrices $\mathcal{T}, \mathcal{S}$.

A tensor $\mathcal{S}$ is called *singular* if the system

$$\mathcal{S} \times \otimes^{d-1}\mathbf{x} = \mathbf{0} \tag{5.3}$$

has a nontrivial solution. Otherwise $\mathcal{S}$ is called nonsingular. It is very easy to give an example of a symmetric nonsingular $\mathcal{S}$ [5]. Let $\mathbf{w}_1,\ldots,\mathbf{w}_m$ be linearly independent in $\mathbb{C}^m$. Then $\mathcal{S}=\sum_{i=1}^m \otimes^d \mathbf{w}_i$ is nonsingular. The set of singular tensors in $\mathbb{C}^{m^{\times d}}$ is a given by the zero set of some multidimensional resultant [7, Chapter 13]. It can be obtained by using the elimination of variables. Let us denote by res $\mathcal{S}$ the multidimensional resultant corresponding to the system (5.3), which is a homogeneous polynomial in the entries of $\mathcal{S}$ of degree $\mu(m,d)=m(d-1)^{m-1}$ [7, Proposition 1.1]. That is, res $\mathcal{S}$ is an irreducible polynomial in $\mathbb{C}^{m^{\times d}}$ such that the system (5.3) as a nonzero solution if and only if res $\mathcal{S}=0$. Furthermore for a general point $\mathcal{S}\in Z(\text{res })$, i.e. the zero set of the polynomial res , the system (5.3) has exactly one simple solution in $\mathbb{P}(\mathbb{C}^m)$. The eigenvalue problem (5.2) consists of two steps. First find all λ satisfying res $(\lambda\mathcal{S}-\mathcal{T})=0$. Clearly res $(\lambda\mathcal{S}-\mathcal{T})$ is a polynomial in λ of degree $\mu(m,d)$ at most. (It is possible that this polynomial in λ is a zero polynomial. This is the case where there exists a nontrivial solution to the system $\mathcal{S}\otimes^{d-1}\mathbf{x}=\mathcal{T}\otimes^{d-1}\mathbf{x}=\mathbf{0}$.) After then one needs to find the nonzero solutions of the system $(\lambda\mathcal{S}-\mathcal{T})\otimes^{d-1}\mathbf{x}=0$ which are viewed as eigenvectors in $\mathbb{P}(\mathbb{C}^m)$. Assume that $\mathcal{S}$ is nonsingular. Then res $(\lambda\mathcal{S}-\mathcal{T})=$ res $(\mathcal{S})\lambda^{\mu(m,d)}+$ polynomial in λ of degree $\mu(m,d)-1$ at most. We show below a result known to the experts, that for general $\mathcal{S},\mathcal{T}$ each eigenvalue λ the system $(\lambda\mathcal{S}-\mathcal{T})\otimes^{d-1}\mathbf{x}=0$ has exactly one corresponding eigenvector in $\mathbb{P}(\mathbb{C}^m)$. We outline a short proof of the following known theorem which basically uses only the existence of the resultant for the system (5.3). For an identity tensor $\mathcal{S}$. i.e. (5.1), see [19].

Theorem 8 *Let $\mathcal{S},\mathcal{T}\in\mathbb{C}^{m^{\times d}}$ and assume that $\mathcal{S}$ is nonsingular. Then* res $(\lambda\mathcal{S}-\mathcal{T})$ *is a polynomial in λ of degree $m(d-1)^{m-1}$. For a generic $\mathcal{S}$ and $\mathcal{T}$ to each eigenvalue λ of the pencil* (5.1) *corresponds one eigenvector in $\mathbb{P}(\mathbb{C}^m)$.*

Proof. Consider the space $\mathbb{P}(\mathbb{C}^2)\times\mathbb{P}(\mathbb{C}^{m^{\times d}}\times\mathbb{C}^{m^{\times d}})\times\mathbb{P}(\mathbb{C}^m)$ with the local coordinates $((u,v),(\mathcal{S},\mathcal{T}),\mathbf{x})$. Consider the system of m-equation homogenous in $(u,v),(\mathcal{S},\mathcal{T}),\mathbf{x}$ given by

$$(u\mathcal{S}-v\mathcal{T})\times\otimes^{d-1}\mathbf{x}=\mathbf{0}. \tag{5.4}$$

The existence of the multidimensional resultant is equivalent to the assumption that the above variety $V(m,d)$ is an irreducible variety of dimension $2m^d-1$ in $\mathbb{P}(\mathbb{C}^2)\times\mathbb{P}(\mathbb{C}^{m^{\times d}}\times\mathbb{C}^{m^{\times d}})\times\mathbb{P}(\mathbb{C}^m)$. So it is enough to find a good point $(\mathcal{S}_0,\mathcal{T}_0)$ such that it has exactly $\mu(m,d)=m(d-1)^{m-1}$ smooth point $((u_i,v_i),(\mathcal{S}_0,\mathcal{T}_0),\mathbf{x}_i)$ in $V(m,d)$.

To do that we assume that $\mathcal{S}_0,\mathcal{T}_0$ are in fact "matrix type" tensors.

$$\mathcal{S}_0\times\otimes^{d-1}\mathbf{x}=A\mathbf{x}^{\circ(d-1)},\ \mathcal{T}_0\times\otimes^{d-1}\mathbf{x}=B\mathbf{x}^{\circ(d-1)},\ A=[a_{ij}],B=b_{ij}]\in\mathbb{C}^{m\times m}. \tag{5.5}$$

to be specific we can choose $A=I$ and B as a cyclic permutation matrix $B(x_1,\ldots,x_m)^\top=(x_2,\ldots,x_m,x_1)^\top$. Then B has m distinct eigenvalues, the $m-th$ roots of unity. $\mathbf{x}$ is an eigenvector of (5.5) if and only if $\mathbf{x}^{\circ(d-1)}$ is an eigenvector of B. Fix an eigenvalue of B. One can fix $x_1=1$. Then we have exactly $(d-1)^{m-1}$ eigenvectors in $\mathbb{P}(\mathbb{C}^m)$ corresponding to each eigenvalue λ of B. So altogether we have $m(d-1)^{m-1}$ distinct eigenvectors. It is left to show that each point $((u_i,v_i),(\mathcal{S}_0,\mathcal{T}_0),\mathbf{x}_i)$ is a simple point of $V(m,d)$. For that we need to show that the Jacobian of the system (5.4) at each point has rank m, the maximal possible rank, at $((u_i,v_i),(\mathcal{S}_0,\mathcal{T}_0),\mathbf{x}_i)$.

For that we assume that $u_i = \lambda_i, v_i = 1, x_1 = 1$. This easily follows from the fact that each eigenvalue of B is a simple eigenvalue. Hence the projection of $V(m,d)$ on $\mathbb{P}(\mathbb{C}^{m^{\times d}} \times \mathbb{C}^{m^{\times d}})$ is $m(d-1)^{m-1}$ valued.

Note that in this example each eigenvalue λ of (5.5) is of multiplicity $(d-1)^{m-1}$. It is left to show that when we consider the pairs $\mathcal{S}_0, \mathcal{T}$ where $\mathcal{T}$ varies in the neighborhood of $\mathcal{T}_0$ we obtain $m(d-1)^{m-1}$ different eigenvalues. Since the Jacobian of the system (5.5) has rank m at each eigenvalue $\lambda_i = \frac{u_i}{v_i}$ and the corresponding eigenvector $\mathbf{x}_i$, one has a simple variation formula for each $\delta\lambda_i$ using the implicit function theorem. Fix $x_1 = 1$ and denote $\mathbf{F}(\mathbf{x}, \lambda, \mathcal{T}) = (F_1, \ldots, F_m) := (\lambda\mathcal{S}_0 - \mathcal{T}) \times \otimes^{d-1}\mathbf{x}$. Thus we have the system of m equations $\mathbf{F}(\mathbf{x}, \lambda) = 0$ in m variables $x_2, \ldots, x_m, \lambda$. We let $\mathcal{T} = \mathcal{T}_0 + t\mathcal{T}_1$ and we want to find the first term of $\lambda_i(t) = \lambda_i + \alpha_i t + O(t^2)$. We also assume that $\mathbf{x}_i(t) = \mathbf{x}_i + t\mathbf{y}_i + O(t^2)$, where $\mathbf{y}_i = (0, y_{2,i}, \ldots, y_{m,i})^\top$. Hence the first order computations yield the equations

$$\begin{aligned} &\mathcal{T}_1 \times \otimes^{d-1}\mathbf{x}_i + \mathcal{T}_0 \times \mathbf{z}_i = \alpha_i \mathcal{S}_0 \times \otimes^{d-1}\mathbf{x}_i + \lambda_i \mathcal{S} \times \mathbf{z}_i, \\ &\mathbf{z}_i = \sum_{j\in[d-1]} \otimes^{j-1}\mathbf{x}_i \otimes \mathbf{y}_i \otimes^{d-1-j} \mathbf{x}_i. \end{aligned} \tag{5.6}$$

Let $\mathbf{w} = (w_1, \ldots, w_m)^\top$ be the left eigenvector of B corresponding to λ_i, i.e. $\mathbf{w}^\top B = \lambda_i \mathbf{w}^\top$ normalized by the condition $\mathbf{w}^\top(\mathcal{S}_0 \times \otimes^{d-1}\mathbf{x}_i) = (\mathcal{S}_0 \times \otimes^{d-1}\mathbf{x}_i) \times \mathbf{w} = 1$. Contracting both sides of (5.6) with the vector $\mathbf{w}$ we obtain

$$\alpha_i = \mathcal{T}_1 \times (\mathbf{w} \otimes (\otimes^{d-1}\mathbf{x}_i)). \tag{5.7}$$

It is straightforward to show that $\alpha_1, \ldots, \alpha_{m(d-1)^{m-1}}$ are pairwise distinct for a general $\mathcal{T}_1$. □

The proof of Theorem 8 yields.

Corollary 9 *Let $\mathcal{T} \in \mathbb{C}^{m^{\times d}})$ be a general tensor. Then the homogeneous eigenvalue problem* (5.1) *has exactly $m(d-1)^{m-1}$ distinct eigenvectors in $\mathbb{P}(\mathbb{C}^m)$ which correspond to distinct eigenvalues.*

We close this section with an heuristic argument which shows that a generic pencil $(\mathcal{S}, \mathcal{T}) \in \mathbb{P}(\mathbb{C}^{m^{\times d}} \times \mathbb{C}^{m^{\times d}})$ has $\mu(m,d) = m(d-1)^{m-1}$ distinct eigenvalues in $\mathbb{P}(\mathbb{C}^m)$. Let $\mathcal{S} \in \mathbb{C}^{m^{\times d}}$ be nonsingular Then $\mathcal{S}$ induces a linear map $\hat{\mathcal{S}}$ from the line bundle $\otimes^{d-1}T(m)$ to the trivial bundle $\mathbb{C}^m$ over $\mathbb{P}(\mathbb{C}^m)$ by $\otimes^{d-1}\mathbf{x} \mapsto \mathcal{S} \times \otimes^{d-1}\mathbf{x}$. Then we have an exact sequence of line bundles

$$0 \to \otimes^{d-1}T(m) \to \mathbb{C}^m \to Q_{m,d} \to 0$$

where $Q_{m,d} = \mathbb{C}^m/(\hat{\mathcal{S}}(\otimes^{d-1}T(m)))$. The Chern polynomial of $Q_{m,d}$ is $1 + \sum_{i=1}^{m-1}(d-1)^i t^i \alpha^i$. A similar computation for finding the number of eigenvectors of (4.1) shows that the number of eigenvalues of (5.2) is the coefficient of t_1^{m-1} in the polynomial $\frac{\hat{t}_1^m - \tilde{t}_1^m}{\hat{t}_1 - \tilde{t}_1}$. Here $\hat{t}_1 = \tilde{t}_1 = (d-1)t_1$. Hence the coefficient of t_1^{m-1} is $\frac{(d-1)^m - (d-1)^m}{(d-1)-(d-1)}$. The calculus interpretation of this formula is the derivative of t^m at $t = d-1$, which gives the value of the coefficient $m(d-1)^{m-1}$.

6 On uniqueness of best approximations

Let $\langle \cdot, \cdot \rangle, \|\cdot\|$ be the standard inner product and the corresponding Euclidean norm on $\mathbb{R}^n$. Let $C \subsetneq \mathbb{R}^n$ be a given nonempty closed set. For each $\mathbf{x} \in \mathbb{R}^n$ we consider the function $\mathrm{dist}(\mathbf{x}, C) := \min\{\|\mathbf{x} - \mathbf{y}\|,\ \mathbf{y} \in C\}$. A point $\mathbf{y}^\star \in C$ is called a best approximation of $\mathbf{x}$ if $\|\mathbf{x} - \mathbf{y}^\star\| = \mathrm{dist}(\mathbf{x}, C)$. The following result is well known and we bring its short proof for completeness.

Lemma 10 *Let $C \subsetneq \mathbb{R}^n$ be a given closed set. Let $\mathbf{U} \subset \mathbb{R}^n$ be a subspace with $\dim \mathbf{U} \in [n]$ and such that $\mathbf{U}$ is not contained in C. Let $d(\mathbf{x}), \mathbf{x} \in \mathbf{U}$ be the restriction of $\mathrm{dist}(\cdot, C)$ to $\mathbf{U}$.*

- *(i) The function $\mathrm{dist}(\cdot, C)$ is Lipschitz with constant constant 1:*

$$|\mathrm{dist}(\mathbf{x}, C) - \mathrm{dist}(\mathbf{z}, C)| \le \|\mathbf{x} - \mathbf{z}\| \text{ for all } \mathbf{x}, \mathbf{z} \in \mathbb{R}^n. \tag{6.1}$$

- *(ii) The function $d(\cdot)$ is differentiable a.e. in $\mathbf{U}$.*

- *(iii) Let $\mathbf{x} \in \mathbf{U} \setminus C$ and assume that $d(\cdot)$ is differentiable at $\mathbf{x}$. Denote the differential as $\partial d(\mathbf{x})$ which is viewed as linear functional on $\mathbf{U}$. Let $\mathbf{y}^\star \in C$ be a best approximation to $\mathbf{x}$. Then*

$$\partial d(\mathbf{x})(\mathbf{u}) = \langle \mathbf{u}, \frac{1}{\mathrm{dist}(\mathbf{x}, C)}(\mathbf{x} - \mathbf{y}^\star)\rangle \text{ for each } \mathbf{u} \in \mathbf{U}. \tag{6.2}$$

  If $\mathbf{z}^\star$ is another best approximation to $\mathbf{x}$ then $\mathbf{z}^\star - \mathbf{y}^\star$ is orthogonal to $\mathbf{U}$.

Proof. Assume that $\mathrm{dist}(\mathbf{x}, C) = \|\mathbf{x} - \mathbf{y}^\star\|$. Then

$$\mathrm{dist}(\mathbf{z}, C) \le \|\mathbf{z} - \mathbf{y}^\star\| \text{ for each } \mathbf{z} \in \mathbb{R}^n. \tag{6.3}$$

Suppose next that $\mathrm{dist}(\mathbf{z}, C) = \|\mathbf{z} - \mathbf{y}\|, \mathbf{y} \in C$. Hence

$$-\|\mathbf{x}-\mathbf{z}\| \le \|\mathbf{x}-\mathbf{y}^\star\| - \|\mathbf{z}-\mathbf{y}^\star\| \le \mathrm{dist}(\mathbf{x}, C) - \mathrm{dist}(\mathbf{z}, C) \le \|\mathbf{x}-\mathbf{y}\| - \|\mathbf{z}-\mathbf{y}\| \le \|\mathbf{x}-\mathbf{z}\|.$$

This proves (6.1) and (i). We get that $d(\cdot)$ is also Lipschitz on $\mathbf{U}$. Rademacher's theorem yields that $d(\cdot)$ is differentiable a.e., so proving (ii). In order to prove (iii), fix $\mathbf{u} \in \mathbf{U}$. Then

$$\mathrm{dist}(\mathbf{x} + t\mathbf{u}, C) = \mathrm{dist}(\mathbf{x}, C) + t\partial d(\mathbf{x})(\mathbf{u}) + to(|t|).$$

(6.3) yields the inequality

$$\mathrm{dist}(\mathbf{x} + t\mathbf{u}, C) \le \|\mathbf{x} + t\mathbf{u} - \mathbf{y}^\star\| = \|\mathbf{x} - \mathbf{y}^*\| + t\langle \mathbf{u}, \frac{1}{\mathrm{dist}(\mathbf{x}, C)}(\mathbf{x} - \mathbf{y}^\star)\rangle + O(t^2).$$

Compare this inequality with the previous equality to deduce that

$$t\partial d(\mathbf{x})(\mathbf{u}) \le t\langle \mathbf{u}, \frac{1}{\mathrm{dist}(\mathbf{x}, C)}(\mathbf{x} - \mathbf{y}^\star)\rangle$$

for all $t \in \mathbb{R}$. This implies (6.2). If $\mathbf{z}^\star$ another best approximation to $\mathbf{x}$ then (6.2) yields that $\mathbf{z}^\star - \mathbf{y}^\star$ is orthogonal to $\mathbf{U}$. $\square$

Corollary 11 *Let $C \subsetneq \mathbb{R}^n$ be a given closed set.*

- *(i) The function* $\mathrm{dist}(\mathbf{x}, C)$ *has differential a.e. in* $\mathbb{R}^n$.
- *(ii) Let* $\mathbf{x} \in \mathbb{R}^n \setminus C$ *and assume that* $\mathrm{dist}(\cdot, C)$ *is differentiable at* $\mathbf{x}$. *Then* $\mathbf{x}$ *has a unique best approximation* $\mathbf{y}(\mathbf{x}) \in C$. *Furthermore*

$$\partial \mathrm{dist}(\mathbf{x}, C)(\mathbf{u}) = \langle \mathbf{u}, \frac{1}{\mathrm{dist}(\mathbf{x}, C)}(\mathbf{x} - \mathbf{y}(\mathbf{x}))\rangle \text{ for each } \mathbf{u} \in \mathbb{R}^n. \tag{6.4}$$

In particular, a.a. $\mathbf{x} \in \mathbb{R}^n$ *have a unique best approximation* $\mathbf{y}(\mathbf{x}) \in C$.

Proof. Choose $\mathbf{U} = \mathbb{R}^n$, so $d(\cdot) = \mathrm{dist}(\cdot, C)$ is differentiable a.e. by Lemma 10 (ii). In order to prove (ii), assume that $\mathbf{y}^\star$ and $\mathbf{z}^\star$ are best approximations of $\mathbf{x}$. Then $\mathbf{z}^\star - \mathbf{y}^\star$ is orthogonal to $\mathbb{R}^n$ by Lemma 10 (iii). So $\mathbf{z}^\star - \mathbf{y}^\star = \mathbf{0}$. Furthermore (6.4) holds. □

7 Best rank one approximations of d-mode tensors

On $\mathbb{C}^{\mathbf{m}}$ define an inner product and its corresponding Hilbert-Schmidt norm $\langle \mathcal{T}, \mathcal{S}\rangle := \mathcal{T} \times \bar{\mathcal{S}}, \|\mathcal{T}\| = \sqrt{\langle \mathcal{T}, \mathcal{T}\rangle}$. We first present some known results of best rank one approximations of real tensors. In this section we assume that $\mathbb{F} = \mathbb{R}$ and $\mathcal{T} \in \mathbb{R}^{\mathbf{m}}$. Let $\mathrm{S}^{m-1} \subset \mathbb{R}^m$ be the $m-1$-dimensional sphere $\|\mathbf{x}\| = 1$. Denote by $\mathrm{S}(\mathbf{m})$ the d-product of the spheres $\mathrm{S}^{m_1-1} \times \ldots \times \mathrm{S}^{m_d-1}$. Let $(\mathbf{x}_1, \ldots, \mathbf{x}_d) \in \mathrm{S}(\mathbf{m})$ and associate with $(\mathbf{x}_1, \ldots, \mathbf{x}_d)$ the d one dimensional subspaces $\mathbf{U}_i = \mathrm{span}(\mathbf{x}_i)$, $i \in [d]$. Note that

$$\| \otimes_{i\in[d]} \mathbf{x}_i\| = \prod_{i\in[d]} \|\mathbf{x}_i\| = 1.$$

The projection $P_{\otimes_{i\in[d]}\mathbf{U}_i}(\mathcal{T})$ of $\mathcal{T}$ onto the one dimensional subspace $\mathbf{U} := \otimes_{i\in[d]}\mathbf{U}_i \subset \otimes_{i\in[d]}\mathbb{R}^{m_i}$, is given by

$$f_{\mathcal{T}}(\mathbf{x}_1, \ldots, \mathbf{x}_d) \otimes_{i\in[d]} \mathbf{x}_i, f_{\mathcal{T}}(\mathbf{x}_1, \ldots, \mathbf{x}_d) := \langle \mathcal{T}, \otimes_{i\in[d]}\mathbf{x}_i\rangle, (\mathbf{x}_1, \ldots, \mathbf{x}_d) \in \mathrm{S}(\mathbf{m}). \tag{7.1}$$

Let $P_{(\otimes_{i\in[d]}\mathbf{U}_i)^\perp}(\mathcal{T})$ be the orthogonal projection of $\mathcal{T}$ onto the orthogonal complement of $\otimes_{i\in[d]}\mathbf{U}_i$. The Pythagorean identity yields

$$\|\mathcal{T}\|^2 = \|P_{\otimes_{i\in[d]}\mathbf{U}_i}(\mathcal{T})\|^2 + \|P_{(\otimes_{i\in[d]}\mathbf{U}_i)^\perp}(\mathcal{T})\|^2. \tag{7.2}$$

With this notation, the best rank one approximation of $\mathcal{T}$ from $\mathrm{S}(\mathbf{m})$ is given by

$$\min_{(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in\mathrm{S}(\mathbf{m})} \min_{a\in\mathbb{R}} \|\mathcal{T} - a \otimes_{i\in[d]} \mathbf{x}_i\|.$$

Observing that

$$\min_{a\in\mathbb{R}} \|\mathcal{T} - a \otimes_{i\in[d]} \mathbf{x}_i\| = \|\mathcal{T} - P_{\otimes_{i\in[d]}\mathbf{U}_i}(\mathcal{T})\| = \|P_{(\otimes_{i\in[d]}\mathbf{U}_i)^\perp}(\mathcal{T})\|,$$

it follows that the best rank one approximation is obtained by the minimization of $\|P_{(\otimes_{i\in[d]}\mathbf{U}_i)^\perp}(\mathcal{T})\|$. In view of (7.2) we deduce that best rank one approximation is obtained by the maximization of $\|P_{\otimes_{i\in[d]}\mathbf{U}_i}(\mathcal{T})\|$ and finally, using (7.1), it follows that the best rank one approximation is given by

$$\sigma_1(\mathcal{T}) := \max_{(\mathbf{x}_1,\ldots,\mathbf{x}_d)\in \mathrm{S}(\mathbf{m})} f_{\mathcal{T}}(\mathbf{x}_1,\ldots,\mathbf{x}_d). \tag{7.3}$$

As in the matrix case $\sigma_1(\mathcal{T})$ is called in [11] the *spectral norm.* Furthermore it is shown in [11] that the computation of $\sigma_1(\mathcal{T})$ in general is NP-hard for $d > 2$.

We will make use of the following result of [14], where we present the proof for completeness.

Lemma 12 *For $\mathcal{T} \in \mathbb{R}^{\mathbf{m}}$, the critical points of $f|_{\mathrm{S}(\mathbf{m})}$, defined in* (7.1)*, are singular value tuples satisfying*

$$\mathcal{T} \times (\otimes_{j\in[d]\setminus\{i\}}\mathbf{x}_j) = \lambda \mathbf{x}_i \quad \textit{for all } i \in [d],\ (\mathbf{x}_1,\ldots,\mathbf{x}_d) \in \mathrm{S}(\mathbf{m}). \tag{7.4}$$

Proof. We need to find the critical points of $\langle \mathcal{T}, \otimes_{j\in[d]}\mathbf{x}_j\rangle$ where $(\mathbf{x}_1,\ldots,\mathbf{x}_d) \in \mathrm{S}(\mathbf{m})$. Using Lagrange multipliers we consider the auxiliary function

$$g(\mathbf{x}_1,\ldots,\mathbf{x}_d) := \langle \mathcal{T}, \otimes_{j\in[d]}\mathbf{x}_j\rangle - \sum_{j\in[d]} \lambda_j \mathbf{x}_j^\top \mathbf{x}_j.$$

The critical points of g then satisfy

$$\mathcal{T} \times (\otimes_{j\in[d]\setminus\{i\}}\mathbf{x}_j) = \lambda_i \mathbf{x}_i, \quad i \in [d],$$

and hence $\langle \mathcal{T}, \otimes_{j\in[d]}\mathbf{x}_j\rangle = \lambda_i \mathbf{x}_i^\top \mathbf{x}_i = \lambda_i$ for all $i \in [d]$ which implies (7.4). □

Observe next that $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ satisfies (7.4) if and only if the vectors $(\pm\mathbf{x}_1,\ldots,\pm\mathbf{x}_d)$ satisfy (7.4). In particular, we could choose the signs in $(\pm\mathbf{x}_1,\ldots,\pm\mathbf{x}_d)$ such that each corresponding λ is nonnegative and then these λ can be interpreted as the singular values of $\mathcal{T}$. The maximal singular value of $\mathcal{T}$ is denoted by $\sigma_1(\mathcal{T})$ and is given by (7.3). Note that to each nonnegative singular value there are at least 2^{d-1} singular vector tuples of the form $(\pm\mathbf{x}_1,\ldots,\pm\mathbf{x}_d)$. So it is more natural to view the singular vector tuples $(\mathbf{x}_1,\ldots,\mathbf{x}_d)$ as points in the real projective Segre variety $\Sigma_{\mathbb{R}}(\mathbf{m})$. Furthermore, the projection of $\mathcal{T}$ on the one dimensional subspace spanned by $\otimes_{i\in[d]}(\pm\mathbf{x}_i)$, where $(\mathbf{x}_1,\ldots,\mathbf{x}_d) \in \mathrm{S}(\mathbf{m})$, is equal to one vector $(\mathcal{T} \times \otimes_{i\in[d]}\mathbf{x}_i) \otimes_{i\in[d]} \mathbf{x}_i$.

Theorem 13 *There exists an variety $V \subset \mathbb{R}^{\mathbf{m}}$ such that for each $\mathcal{T} \in \mathbb{R}^{\mathbf{m}} \setminus V$ the best rank one approximation is unique.*

Proof. Theorem 4 implies the existence of a variety $V_0 \subset \mathbb{C}^{\mathbf{m}}$ such that each $\mathcal{T} \in \mathbb{C}^{\mathbf{m}} \setminus V_0$ has exactly $c(\mathbf{m})$ distinct singular value tuples. Lemma 12 implies for any $\mathcal{T} \in \mathbb{R}^{\mathbf{m}}$ the best rank approximation is of the form $\lambda_p \otimes_{j\in[d]} \mathbf{x}_{j,p}$, Here $(\mathbf{x}_{1,p},\ldots,\mathbf{x}_{d,p}) \in \mathrm{S}(\mathbf{m})$ satisfies (7.4) with $\lambda = \lambda_p \geq 0$ and $\mathbf{x}_j = \mathbf{x}_{j,p}, j \in [d]$. Furthermore $\lambda_p = \sigma_1(\mathcal{T})$, where $\sigma_1(\mathcal{T})$ is the maximum of all λ_p. Assume that $\mathcal{T} \in \mathbb{R}^{\mathbf{m}} \setminus V_0$. Then the number of rank one tensors $\otimes_{j\in[d]}\mathbf{x}_{j,p}$ is at most $c(\mathbf{m})$. Hence there exists a real subvariety $V \subset \mathbb{R}^{\mathbf{m}}$, which contains $V_0 \cap \mathbb{R}^{\mathbf{m}}$ such there are exactly $k \geq 1$ different rank one tensor $\otimes_{j\in[d]}\mathbf{x}_{j,p}$ such that $\lambda_p = \sigma_1(\mathcal{T})$. This implies that that for each $\mathcal{T} \in \mathbb{R}^{\mathbf{m}} \setminus V$ we have exactly k best rank one approximation. Recall now Corollary 11 for C being the variety of all tensors of rank one plus the zero tensor. This Corollary yields that for almost all $\mathcal{T}$ the best rank one approximation is unique. Hence $k = 1$. $\square$

Let $\boldsymbol{\omega} = (\omega_1, \ldots, \omega_p)$ be a partition of d. For $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ it is natural to consider a best rank one approximation to $\mathcal{T}$ of the form $\pm \prod_{i\in[p]} \otimes^{\omega_i} \mathbf{x}_i$ where $(\mathbf{x}_1, \ldots, \mathbf{x}_p) \in \Sigma_{\mathbb{R}}(\mathbf{m}')$. We call such an approximation a best $\boldsymbol{\omega}$-symmetric rank one approximation. (The factor $\pm$ is needed only if each ω_i is even.) As in the case $\mathcal{T} \in \mathbb{R}^{\mathbf{m}}$ a best $\boldsymbol{\omega}$-symmetric rank one approximation of $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ is a solution to the following maximum problem.

$$\max_{(\mathbf{x}_1,\ldots,\mathbf{x}_p)\in \mathrm{S}(\mathbf{m}')} |\mathcal{T} \times \otimes_{i\in[p]} \otimes^{\omega_i} \mathbf{x}_i|. \tag{7.5}$$

As before, the critical points of the functions $\pm\mathcal{T} \times \otimes_{i\in[p]} \otimes^{\omega_i} \mathbf{x}_i$ on $\mathrm{S}(\mathbf{m}')$ satisfy

$$\mathcal{T} \times \otimes_{j\in[p]} \otimes^{\omega_j - \delta_{ji}} \mathbf{x}_j = \lambda \mathbf{x}_i, \quad i \in [p], \quad (\mathbf{x}_1, \ldots, \mathbf{x}_p) \in \mathrm{S}(\mathbf{m}'). \tag{7.6}$$

The best $\boldsymbol{\omega}$-symmetric rank one approximation corresponds to all λ for which $|\lambda|$ has a maximal possible value. Combine the arguments of the proof of Theorem 13 with Theorem 5 to obtain.

Proposition 14 *There exists a variety $U \subset \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ such that for each $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \setminus U$ the best rank one $\boldsymbol{\omega}$-symmetric approximation is unique.*

Assume that $\otimes_{j\in[d]} \mathbf{y}_j \in \mathbb{R}^{\mathbf{m}(\boldsymbol{\omega})}$ is a best rank one approximation to a tensor $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$. It is not obvious a priori that $\otimes_{j\in[d]} \mathbf{y}_j$ is $\boldsymbol{\omega}$-symmetric. However, the following result is obvious.

$$\begin{array}{l} \otimes_{j\in[d]} \mathbf{y}_{\sigma(j)} \text{ is best rank one approximation of } \mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \\ \text{for each permutation } \sigma : [d] \to [d] \text{ which preserves } \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}). \end{array} \tag{7.7}$$

Lemma 15 *For a.a. $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ there exists a unique rank one tensor $\otimes_{j\in[d]} \mathbf{y}_j \in \mathbb{R}^{\mathbf{m}(\boldsymbol{\omega})}$ such that all best rank one approximations of $\mathcal{T}$ are of the form* (7.7).

To prove this lemma we need an auxiliary lemma.

Lemma 16 *Let $\otimes_{j\in[d]} \mathbf{x}_j, \otimes_{j\in[d]} \mathbf{y}_j \in \mathbb{R}^{n^{\times d}}$. Assume that*

$$\langle \otimes_{j\in d} \mathbf{x}_j, \otimes^d \mathbf{u} \rangle = \langle \otimes_{j\in d} \mathbf{y}_j, \otimes^d \mathbf{u} \rangle \ \forall \mathbf{u} \in \mathbb{R}^n. \tag{7.8}$$

Then there exists a permutation $\sigma : [d] \to [d]$ such that $\otimes_{j\in[d]} \mathbf{y}_j = \otimes_{j\in[d]} \mathbf{x}_{\sigma(j)}$.

Proof. Note that the condition (7.8) is equivalent to the equality

$$\prod_{j\in[d]} \mathbf{u}^\top \mathbf{x}_j = \prod_{j\in[d]} \mathbf{u}^\top \mathbf{y}_j \ \forall \mathbf{u} \in \mathbb{R}^n. \tag{7.9}$$

If $\otimes_{j\in d} \mathbf{x}_j = 0$ then $\prod_{j\in[d]} \mathbf{u}^\top \mathbf{y}_j = 0$ for all $\mathbf{u}$. Hence $\mathbf{y}_j = \mathbf{0}$ for some j, so $\otimes_{j\in[d]} \mathbf{y}_j = \otimes_{j\in[d]} \mathbf{x}_j = 0$. So we assume that $\otimes_{j\in[d]} \mathbf{x}_j, \otimes_{j\in[d]} \mathbf{y}_j$ are both nonzero.

We now prove the lemma by induction. For $d = 1$ the lemma is trivial. So assume that the lemma hold of $d = k$. Let $d = k + 1$. Assume that $\mathbf{u} \in \mathrm{span}(\mathbf{x}_{k+1})^\perp$. Then (7.9) yields that $\prod_{j\in[d]} \mathbf{u}^\top \mathbf{y}_j = 0$. Hence $\mathrm{span}(\mathbf{x}_{k+1})^\perp \subset \cup_{j\in[k+1]}\mathrm{span}(\mathbf{y}_j)$. Therefore there exists $j \in [k + 1]$ such that $\mathrm{span}(\mathbf{x}_{k+1})^\perp = \mathrm{span}(\mathbf{y}_j)^\perp$. So $\mathbf{y}_j = t\mathbf{x}_{k+1}$ for some $t \in \mathbb{R} \setminus \{0\}$. Hence there exist $\mathbf{z}_1, \ldots, \mathbf{z}_{d+1} \in \mathbb{R}^n$ and a permutation $\sigma' : [k + 1] \to [k + 1]$ such that $\otimes_{j\in[k+1]}\mathbf{z}_{\sigma'(j)} = \otimes_{j\in[k+1]}\mathbf{y}_j$ where $\mathbf{z}_{k+1} = \mathbf{x}_{k+1}$. Hence $\otimes_{j\in[k+1]}\mathbf{x}_j$ and $\otimes_{j\in[k+1]}\mathbf{z}_j$ satisfy (7.9). Therefore $\otimes_{j\in[k]}\mathbf{x}_j$ and $\otimes_{j\in[k]}\mathbf{z}_j$ satisfy (7.9). Use the induction hypotesis to deduce the lemma. □

Proof of Lemma 15. We use Corollary 11 as follows. Let $\mathbb{R}^n = \mathbb{R}^{\mathbf{m}(\boldsymbol{\omega})}$ and assume that C is the set of all rank one tensors plus the zero tensor. We let $\mathbf{U} := \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$. Assume that $d(\cdot)$ is differentiable at $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \setminus C$. Suppose that $\otimes_{j\in[d]}\mathbf{y}_j, \otimes_{j\in[d]}\mathbf{z}_j$ are best rank approximations of $\mathcal{T}$. So

$$\sigma_1(\mathcal{T}) = \| \otimes_{j\in[d]} \mathbf{y}_j\| = \prod_{j\in[d]} \|\mathbf{y}_j\| = \| \otimes_{j\in[d]} \mathbf{z}_j\| = \prod_{j\in[d]} \|\mathbf{z}_j\| > 0.$$

Without loss of generality we may assume that

$$\|\mathbf{y}_j\| = \|\mathbf{z}_j\| = \sigma_1(\mathcal{T})^{\frac{1}{d}} \ \forall j \in [d]. \tag{7.10}$$

Corollary 11 yields that

$$\langle \otimes_{i\in[p]} \otimes^{\omega_i} \mathbf{u}_i, \otimes_{j\in[d]}\mathbf{y}_j - \otimes_{j\in[d]}\mathbf{z}_j\rangle = 0 \ \forall \mathbf{u}_i \in \mathbb{R}^{m'_i} \ i \in [p].$$

The above equality is equivalent to

$$\prod_{i\in[p]} \prod_{j_i\in[\omega_i]} \mathbf{u}_i^\top \mathbf{y}_{\alpha_i+j_i} = \prod_{i\in[p]} \prod_{j_i\in[\omega_i]} \mathbf{u}_i^\top \mathbf{z}_{\alpha_i+j_i}, \quad \forall \mathbf{u}_i \in \mathbb{R}^{m'_i},\ i \in [p], \tag{7.11}$$

where $\omega_0 = 0$ and $\alpha_i = \sum_{k=0}^{i-1} \omega_k$ for all $i \in [p]$.

Suppose first that $p = 1$, i.e. $\mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ is the set of all symmetric tensors in $\mathbb{R}^{m_1^{\times\omega_1}}$. (Note that $d = \omega_1$.) Then Lemma 16 and (7.11) yields that $\otimes_{j\in[d]}\mathbf{z}_j = \otimes_{j\in[d]}\mathbf{y}_{\sigma(j)}$ for some permutation $\sigma : [d] \to [d]$. This proves our lemma for $p = 1$.

Assume now that $p > 1$. Fix $k \in [p]$. Fix $\mathbf{u}_i \in i \in [p] \setminus \{k\}$. Let

$$s_k := \prod_{i\in[p]\setminus\{k\}} \prod_{l_j\in[\omega_j]} \mathbf{u}_{l_j}^\top \mathbf{y}_{\alpha_j+l_j}, \quad t_k := \prod_{i\in[p]\setminus\{k\}} \prod_{l_j\in[\omega_j]} \mathbf{u}_{l_j}^\top \mathbf{z}_{\alpha_j+l_j}.$$

Assume that $s_k \neq 0$. Then the two rank one tensors $s_k \otimes_{l_k\in[\omega_k]} \mathbf{y}_{\alpha_k+l_k}, t_k \otimes_{l_k\in[\omega_k]} \mathbf{z}_{\alpha_k+l_k} \in \mathbb{R}^{(m'_k)^{\times\omega_k}}$ satisfy the assumptions of Lemma 16. Hence there exists a permutation $\sigma_k : [\omega_k] \to [\omega_k]$ such that $t_k \otimes_{l_k\in[\omega_k]} \mathbf{z}_{\alpha_k+l_k} = s_k \otimes_{l_k\in[\omega_k]} \mathbf{y}_{\alpha_k+\sigma_k(l_k)}$. In view of (7.10) we deduce the equality $\otimes_{l_k\in[\omega_k]}\mathbf{z}_{\alpha_k+l_k} = \pm \otimes_{l_k\in[\omega_k]} \mathbf{y}_{\alpha_k+\sigma_k(l_k)}$. Hence there exists $\omega : [d] \to [d]$ which leaves invariant each set $[\alpha_{j+1}]$ for $j \in [p - 1]$ such that $\otimes_{j\in[d]}\mathbf{z}_j = \pm \times_{j\in[d]} \mathbf{y}_{\sigma(j)}$. As $\otimes_{j\in[d]}\mathbf{z}_j$ and $\otimes_{j\in[d]}\mathbf{y}_j$ are best rank one approximation to $\mathcal{T}$ we deduce that $\otimes_{j\in[d]}\mathbf{z}_j = \otimes_{j\in[d]}\mathbf{y}_{\sigma(j)}$. □

Recall a recent result of the first author which claims that each $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ has a best rank one approximation which is $\boldsymbol{\omega}$-symmetric [5, Theorem 1]. For symmetric tensors this theorem is [3, Theorem 4.1]. Use Lemma 15, Theorem 5, [5, Theorem 1] and arguments analogous to the proof of Theorem 13 to deduce.

Theorem 17 *Each $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ has a best rank one approximation which is $\boldsymbol{\omega}$-symmetric. Furthermore, there exists a variety $W \subset \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ such that for each $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \setminus W$ the best rank one approximation is unique and $\boldsymbol{\omega}$-symmetric.*

We now give an independent proof of this theorem which does not use the results in [5, 3].

Proof of Theorem 17. We first show the result in [3, 5] that for each symmetric tensor $\mathcal{T} \in \mathrm{S}^d(\mathbb{R}^n)$ there exists a best rank one approximation which is also symmetric, i.e. of the form $\pm \otimes^d \mathbf{y}$. We prove that by induction on d. For $d = 2$, i.e. $\mathcal{T}$ is a symmetric matrix, this is a well known result which can be deduced from the singular value decomposition of real symmetric tensors, i.e. [8, 5]. Assume that this claim holds for any $d = N \geq 2$ and suppose that $d = N + 1$. Let $\mathcal{T} \in \mathrm{S}^{N+1}(\mathbb{R}^n)$ satisfy the assumption of Lemma 15. That is, there exist $N+1$ vectors $\mathbf{y}_1, \ldots, \mathbf{y}_{N+1}$ of length one and $a \in \mathbb{R}$ such that all best rank one approximations of $\mathcal{T}$ are of the form $a \otimes_{j\in[N+1]} \mathbf{y}_{\sigma(j)}$, for each permutation $\sigma : [N+1] \to [N+1]$. ($a = \mathcal{T} \times \otimes_{j\in[N+1]} \mathbf{y}_{\sigma(j)}$.) We claim that $\mathcal{S} := a \otimes_{j\in[N+1]} \mathbf{y}_j$ is symmetric. Assume to the contrary that $\mathcal{S}$ is not symmetric. So $a \neq 0$. Consider the symmetric tensor $\mathcal{T}_1 := \mathcal{T} \times \mathbf{y}_{N+1} \in \mathrm{S}^N(\mathbb{R}^n)$. By induction hypothesis there exists a unit vector $\mathbf{x} \in \mathbb{R}^n$ such that $b \otimes^N \mathbf{x}$ is a best rank one approximation of $\mathcal{S}$. Hence $b(\otimes^N \mathbf{x}) \otimes \mathbf{y}_{N+1}$ is a best rank one approximation of $\mathcal{T}$. Therefore we can assume that $\mathbf{y}_i = \mathbf{x}$ for $i \in [N]$. Since $\mathcal{S}$ was not symmetric tensor $\mathbf{y}_N \neq \pm\mathbf{x}$. Let $\mathcal{S}' := \mathcal{T} \times \mathbf{x}$. Repeat the arguments for $\mathcal{S}'$ to deduce that there exists a unit vector $\mathbf{x}' \in \mathbb{R}^n$ such that a best rank approximation of $\mathcal{T}$ of the form $b' \otimes^N \mathbf{x}' \otimes \mathbf{x}$, where $\mathbf{x}' \neq \pm\mathbf{x}$. This contradicts Lemma 15. Hence $\mathcal{T}$ has a unique best rank one approximation $\mathcal{S}$ which is symmetric.

As the set of symmetric tensors satisfying the assumptions of Lemma 15 is dense in $\mathrm{S}^{N+1}(\mathbb{R}^n)$ it follows that each $\mathcal{T} \in \mathrm{S}^{N+1}(\mathbb{R}^n)$ has a best rank one approximation which is symmetric.

Assume now the general case that $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$, where $\omega = (\omega_1, \ldots, \omega_p)$ and $p > 1$. Suppose first that $\mathcal{T}$ satisfies the assumptions of Lemma 15. As in the symmetric case we deduce that a best rank approximation of $\mathcal{T}$ is ω-symmetric and unique. Since the set of such $\mathcal{T}$ is dense in $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$, the arguments of the proof of Theorem 13 yield that there exists a variety $W \subset \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R})$ such that for each $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \setminus W$ the best rank one approximation is unique and $\boldsymbol{\omega}$-symmetric. Let $\mathcal{T} \in W$. Since $\mathcal{T}$ is a limit of tensors in $\mathrm{S}^{\boldsymbol{\omega}}(\mathbb{R}) \setminus W$ it follows that a best rank one approximation of $\mathcal{T}$ can be chosen to be ω-symmetric. □

8 Best rank-$(r_1, \ldots, r_d)$ approximation

Assume that $\mathbb{F}$ is a field and $\mathbf{m} = (m_1, \ldots, m_d) \in \mathbb{N}^d$. Let $M = \prod_{j\in d} m_j$, $M_i = \frac{M}{m_i}$ and $\mathbf{m}_i = (m_1, \ldots, m_{i-1}, m_{i+1}, \ldots, m_d) \in \mathbb{N}^{d-1}$ for $i \in [d]$. Assume that $\mathcal{T} = [t_{i_1,\ldots,i_d}] \in \mathbb{F}^{\mathbf{m}}$. Denote by $T_i \in \mathbb{F}^{m_i \times M_i}$ the unfolded matrix of the tensor $\mathcal{T}$ in the mode i. That is, let $\mathcal{T}_{j,k} \in \mathbb{F}^{\mathbf{m}_k}$ be the following $d-1$ mode tensor. Its entries are $[t_{i_1,\ldots,i_{k-1},j,i_{k+1},\ldots,i_d}]$ for $i_p \in [m_p], p \in [d] \setminus \{k\}$. So $j \in [m_k]$. Then the row j of T_i is a tensor $\mathcal{T}_{j,i}$ viewed as a vector in $\mathbb{F}^{\mathbf{m}_i}$. Then $\mathrm{rank}_i\mathcal{T}$ is the rank of the matrix T_i. T_i can be seen as the matrix of the contraction map $\otimes_{j\in[d]\setminus\{i\}}(\mathbb{F}^\vee)^{m_j} \to \mathbb{F}^{m_i}$ for $i \in [d]$. Clearly,

$$\mathrm{rank}_i\mathcal{T} \leq \min(m_i, M_i) \quad i \in [d]. \tag{8.1}$$

Carlini and Kleppe characterized the possible r_i occurring as in the following Theorem.

Theorem 18 ([1], Theorem 7) *Suppose that* $r_i \in [m_i]$ *for* $i \in [d]$. *Then there exists* $\mathcal{T} \in \mathbb{F}^{\mathbf{m}}$ *such that* $\mathrm{rank}_i \mathcal{T} = r_i$ *for* $i \in [d]$ *if and only if*

$$r_i^2 \le \prod_{j \in [d]} r_j \quad \textit{for each } i \in [d]. \tag{8.2}$$

We show a related argument working on any infinite field. For each i let f_i be one minor of T_i of order $\min(m_i, M_i)$. Let $f = \prod_{i\in[d]} f_i$, which is a nonzero polynomial in the entries of $\mathcal{T} = [t_{i_1,\dots,i_d}]$. Let $V(\mathbf{m}) \subset \mathbb{F}^{\mathbf{m}}$ be the zero set of f.

Theorem 19 *Let* $\mathbf{m} \in \mathbb{N}^d$ *and assume that* $V(\mathbf{m}) \subset \mathbb{F}^{\mathbf{m}}$ *is defined as above. Then for each* $\mathcal{T} \in \mathbb{F}^{\mathbf{m}} \setminus V(\mathbf{m})$ *the following equality holds.*

$$\mathrm{rank}_i \mathcal{T} = \min(m_i, M_i) \textit{ for } i \in [d]. \tag{8.3}$$

In particular for $\mathbb{F}$ *being a infinite field, a general tensor* $\mathcal{T} \in \mathbb{F}^{\mathbf{m}}$ *satisfies* (8.3).

Proof. Suppose first that $m_i \le M_i$. We claim that the m_i tensors $\mathcal{T}_{1,i}, \dots, \mathcal{T}_{m_i,i}$ are linearly independent. Suppose not. Then any $m_i \times m_i$ minor of T_i is zero. This contradicts the assumption that the $\mathcal{T} \in \mathbb{F}^{\mathbf{m}} \setminus V(\mathbf{m})$. Hence $\mathrm{rank}_i \mathcal{T} = m_i$. Suppose that $m_i > M_i$. Let $\mathcal{T}_{k_1,i}, \dots, \mathcal{T}_{k_{M_i},i}$ be the M_i tensors which contribute to the minor f_i. Since $f_i(\mathcal{T}) \ne 0$ we deduce that $\mathcal{T}_{k_1,i}, \dots, \mathcal{T}_{k_{M_i},i}$ are linearly independent. Hence $\mathrm{rank}_i \mathcal{T}_i = M_i$ for each $i \in [d]$. Since f is a nonzero polynomial, for an infinite field $\mathbb{F}$ $V(\mathbf{m})$ is a proper closed subset of $\mathbb{F}^{\mathbf{m}}$ in the Zariski topology. Hence (8.3) holds for a general tensor. □

On infinite fields, Theorem 18 can be proved as a consequence of Theorem 19. Indeed, let $\mathbf{r} = (r_1, \dots, r_d) \in \mathbb{N}^d$ and assume that (8.2) holds. Choose a general $\mathcal{T}' = [t'_{j_1,\dots,j_d}] \in \mathbb{F}^{\mathbf{r}}$. So $\mathrm{rank}_i \mathcal{T}' = r_i, i \in [d]$. Extend $\mathcal{T}'$ to $\mathcal{T} = [t_{i_1,\dots,i_d}] \in \mathbb{F}^{\mathbf{m}}$ by adding zero entries. I.e. $t_{j_1,\dots,j_d} = t'_{j_1,\dots,j_d}$ for $j_i \in [r_i], i \in [d]$, and all other entries of $\mathcal{T}$ are zero. Then $\mathrm{rank}_i \mathcal{T} = r_i, i \in [d]$.

In what follows we assume that $\mathbb{F} = \mathbb{R}$. Observe that the set of tensors having rank-$(r_1, \dots, r_d)$ contains in the closure exactly all tensors of rank-$(a_1, \dots, a_d)$ with $a_i \le r_i$. This closure is an algebraic variety, defined as the zero set of all the minors of order $r_i + 1$ of T_i for $i \in [d]$. We denote it by $C_{\mathbf{r}}$. Note that having rank $(1, \dots, 1)$ is equivalent to have rank 1.

Clearly $C_{\mathbf{r}}$ is a closed set in $\mathbb{R}^{\mathbf{m}}$. The best $\mathbf{r}$-rank approximation of $\mathcal{T}$ is the closest tensor in $C_{\mathbf{r}}$ to $\mathcal{T}$ in L_2 norm [4]. Corollary 11 yields.

Theorem 20 *Let* $\mathbf{m} = (m_1, \dots, m_d), \mathbf{r} = (r_1, \dots, r_d)$ *where* $r_i \in [m_i]$ *for* $i \in [d]$ *and they satisfy (8.2). Then almost all* $\mathcal{T} \in \mathbb{R}^{\mathbf{m}}$ *have a unique best* $\mathbf{r}$*-rank approximation.*

Let $\boldsymbol{\omega} = (\omega_1, \dots, \omega_p)$ be a partition of d, $\mathbf{m}' = (m'_1, \dots, m'_p)$ and assume that $\mathbf{m}(\boldsymbol{\omega})$ is defined by (4.4). Assume that $\mathbf{r}' = (r'_1, \dots, r'_p)$, where $r'_i \in [m'_i]$ for $i \in [p]$. Let $\mathbf{r}(\boldsymbol{\omega}) = (\underbrace{r'_1, \dots, r'_1}_{\omega_1}, \dots, \underbrace{r'_p, \dots, r'_p}_{\omega_p})$.

Let $C'_{\mathbf{r}'} = C_{\mathbf{r}(\boldsymbol{\omega})} \cap \mathrm{S}^{\boldsymbol{\omega}}$. Clearly, $C'_{\mathbf{r}'}$ is a closed set, consisting of ω-symmetric tensors in $\mathbb{R}^{\mathbf{m}(\boldsymbol{\omega})}$ having rank $\mathbf{r}(\boldsymbol{\omega})$.

Let $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$. Then a best ω symmetric $\mathbf{r}(\boldsymbol{\omega})$-rank approximation of $\mathcal{T}$ is the closest tensor in $C'_{\mathbf{r}'}$ to $\mathcal{T}$. Corollary 11 yields.

Theorem 21 *Let* $\boldsymbol{\omega} = (\omega_1, \ldots, \omega_p)$ *be a partition of* d. *Assume that* $\mathbf{m}' = (m'_1, \ldots, m'_p), \mathbf{r}' = (r'_1, \ldots, r'_p), r'_i \in [m'_i], i \in [p]$ *and that* $\mathbf{m}(\boldsymbol{\omega})$ *satisfies (8.2). Then almost all* $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$ *have a unique best* ω*-symmetric* $\mathbf{r}(\boldsymbol{\omega})$*-rank approximation.*

We close our paper with the following problem. Let $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$. Does $\mathcal{T}$ have a best $\mathbf{r}(\boldsymbol{\omega})$-rank approximation which is ω-symmetric? If the answer is yes, is a best $\mathbf{r}(\boldsymbol{\omega})$-rank approximation unique for almost all $\mathcal{T} \in \mathrm{S}^{\boldsymbol{\omega}}$? In the previous section we showed that for $\mathbf{r}(\omega) = (1, \ldots, 1)$ the answers to these problems are yes.

References

[1] E. Carlini and J. Kleppe, Ranks derived from multilinear maps, *Journal of Pure and Applied Algebra*, 215(2011), 1999–2004.

[2] D. Cartwright, B. Sturmfels, The number of eigenvectors of a tensor, to appear in Linear Algebra and its Applications, arXiv:1004.4953

[3] B. Chen, S. He, Z. Li, and S, Zhang, Maximum block improvement and polynomial optimization, *SIAM J. Optimization*, 22 (2012), 87107.

[4] L. de Lathauwer, B. de Moor and J. Vandewalle, On the best rank–1 and rank–$(R_1, \ldots, R_N)$ approximation of higher-order tensors, *SIAM J. Matrix Anal. Appl.* 21 (2000), 1324–1342.

[5] S. Friedland. Best rank one approximation of real symmetric tensors can be chosen symmetric, to appear in Special Issue "Tensor Theory", *Frontiers of Mathematics in China*, Springer, arXiv:1110.5689 .

[6] W. Fulton, Intersection Theory, Springer, Berlin 1984

[7] I. M. Gelfand, M. M. Kapranov, A. V. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants* Birkhäuser, Boston, 1994.

[8] G.H. Golub and C.F. Van Loan. *Matrix Computations.* John Hopkins Univ. Press, Baltimore, Md, USA, 3rd Ed., 1996.

[9] P. Griffiths and J. Harris, *Priniciples of Algebraic Geometry*, Wiley 1978.

[10] R. Hartshorne, Algebraic Geometry, Graduate Texts in Mathematics 52, Springer, 1977, New York

[11] C.J. Hillar and L.-H. Lim. Most tensor problems are NP hard, arXiv:0911.1393.

[12] F. Hirzebruch, *Topological Methods in Algebraic Geometry*, Grundlehren der math. Wissenschaften, vol. 131, Springer, 1966.

[13] S. Kobayashi, *Differential Geometry of Complex Vector Bundles*, Princeton University Press 1987.

[14] L.-H. Lim. Singular values and eigenvalues of tensors: a variational approach. *Proc. IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing* (CAMSAP '05), 1 (2005), 129-132.

[15] L. Lyusternik, and L. Shnirel'man, Topological methods in variational problems and their application to the differential geometry of surfaces. (Russian) Uspehi Matem. Nauk (N.S.) 2, (1947). no. 1(17), 166-217.

[16] C. Massri, Algorithm to find a maximum of a multilinear map over a product of spheres, arXiv:1110.6217

[17] G. Ni, L. Qi, F. Wang,Y. Wang, The degree of the E-characteristic polynomial of an even order tensor, *J. Math. Anal. Appl.* 329(2007), n.2, 1218-1229

[18] L. Oeding, G. Ottaviani, Eigenvectors of tensors and algorithms for Waring decomposition, to appear in Journal of Symbolic Computation, arXiv: 1103.0203

[19] L. Qi, Eigenvalues of a real supersymmetric tensor, *J. Symbolic Comput.* 40 (2005) 1302-1324.

[20] L. Qi: Eigenvalues and invariants of tensors, J. Math. Anal. Appl. 325 (2007) 1363–1377.

[21] X. Zhang, C. Ling, L. Qi, The best rank-1 approximation of a symmetric tensor and related spherical optimization problems, preprint 2012